Information For the Use of
Thiocolchicoside (8mg) + Aceclofenac (100mg) + Paracetamol (325mg) Tablets

Aceflick Th8 Max Tablet

ACEFLICK TH8 MAX TABLET

Thiocolchicoside (8 mg) + Aceclofenac (100 mg) + Paracetamol (325 mg)

1. Composition

Each film-coated tablet contains:
Thiocolchicoside 8 mg + Aceclofenac 100 mg + Paracetamol 325 mg
Excipients q.s.

2. Pharmacological Action

Thiocolchicoside is a centrally acting muscle relaxant with antispasmodic properties. It helps reduce painful skeletal-muscle spasm.

Aceclofenac is a non-steroidal anti-inflammatory drug (NSAID) that inhibits cyclooxygenase-mediated prostaglandin synthesis, thereby reducing pain and inflammation.

Paracetamol provides analgesic and antipyretic effects and contributes to overall pain relief.

The combination provides **muscle relaxation, analgesic and anti-inflammatory effects** in painful musculoskeletal conditions associated with muscle spasm.

3. Indications

Aceflick TH8 Max Tablet may be indicated for short-term symptomatic treatment of painful musculoskeletal conditions associated with muscle spasm, including:

- **Acute low back pain**
- Painful muscle spasm
- Sprains and strains
- Musculoskeletal pain with inflammation
- Other acute painful conditions associated with skeletal-muscle spasm

4. Dosage and Administration

Adults: The usual dose is **one tablet once or twice daily**, or as directed by the physician.

The tablet should be swallowed whole with sufficient water, preferably **after food**.

The treatment should generally be limited to the shortest duration necessary. **Thiocolchicoside should not be used for prolonged treatment**, and the total daily dose should remain within the approved prescribing limits.

Do not take additional medicines containing **paracetamol or other NSAIDs** unless specifically advised by the physician.

5. Precautions

- Use for the **shortest possible duration** at the recommended dose.
- Thiocolchicoside should not be used during **pregnancy or breastfeeding** unless specifically prescribed where permitted by applicable regulations.
- Use cautiously in patients with **hepatic or renal impairment**.
- Patients with a history of peptic ulcer, gastrointestinal bleeding, hypertension, heart failure or cardiovascular disease should be evaluated before NSAID therapy.
- Avoid excessive alcohol consumption because of increased risk of liver injury with paracetamol and gastrointestinal/hepatic adverse effects.
- Thiocolchicoside may cause drowsiness or dizziness; caution is advised while driving or operating machinery.
- Discontinue and seek medical attention if gastrointestinal bleeding, severe skin reactions, allergic reactions or symptoms of liver injury occur.

6. Contraindications

The combination is contraindicated in patients with:

- Known hypersensitivity to **thiocolchicoside, aceclofenac, paracetamol, NSAIDs**, or any component of the formulation.
- Active or recurrent **peptic ulceration or gastrointestinal bleeding**.
- Severe hepatic or renal impairment, where contraindicated by the approved product information.
- Severe heart failure or established NSAID-related contraindications.
- **Pregnancy and breastfeeding** for thiocolchicoside-containing products, according to applicable prescribing restrictions.
- History of bronchospasm, urticaria or allergic reactions precipitated by aspirin or other NSAIDs.

7. Adverse Reactions

Possible adverse reactions include:

- Nausea, vomiting and indigestion
- Abdominal pain or diarrhoea
- Gastritis
- Dizziness or drowsiness
- Headache
- Increased liver enzymes

NSAIDs may rarely cause **gastrointestinal ulceration/bleeding, renal impairment, fluid retention, cardiovascular events and serious hypersensitivity or skin reactions**.

Paracetamol overdose may cause **severe liver injury**.

8. Drug Interactions

Important interactions may occur with:

- **Other NSAIDs or aspirin**, increasing the risk of gastrointestinal and renal adverse effects.
- Other medicines containing **paracetamol**, increasing the risk of overdose and hepatotoxicity.
- Anticoagulants such as **warfarin**, which may increase bleeding risk.
- Corticosteroids, antiplatelet drugs and SSRIs, which may increase gastrointestinal bleeding risk.
- ACE inhibitors, ARBs and diuretics, which may have reduced antihypertensive effect and increased renal risk when combined with NSAIDs.
- Methotrexate, lithium, cyclosporine and other medicines requiring careful monitoring.

Inform the physician about all prescription, OTC and herbal medicines being taken.

9. Storage

Store below **30°C**, protected from light and moisture.

Keep out of reach of children.

10. Note / Important

Aceflick TH8 Max is intended for **short-term symptomatic relief of painful musculoskeletal conditions associated with muscle spasm**.

Do not exceed the prescribed dose. Avoid taking additional paracetamol-containing or NSAID-containing medicines without medical advice.

Because thiocolchicoside has specific safety restrictions, including **dose and duration limitations and important reproductive safety precautions**, use should strictly follow the approved prescribing information.

For use under medical supervision.

Keep out of reach of children.

Manufactured in India for:

Cafoli™
LIFECARE

Cafoli Lifecare Pvt. Ltd.
(An ISO 9001: 2015 Certified Co.)
Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113
Corporate Office :
Plot no. 307, Industrial Area Phase-2,
Panchkula, Haryana -134113
TM: Trademark Applied for