Information For the Use of
Aceclofenac (100mg) + Tizanidine (2mg) Tablets

Aceflick TZ Tablet

ACEFLICK-TZ TABLET

Aceclofenac (100 mg) + Tizanidine (2 mg)

Composition

Each tablet contains:

- **Aceclofenac – 100 mg**
- **Tizanidine – 2 mg**

Pharmacological Action

Aceclofenac is a non-steroidal anti-inflammatory drug (NSAID) that provides analgesic and anti-inflammatory activity, while tizanidine is a centrally acting skeletal muscle relaxant.

It functions by:

- **Analgesic & Anti-inflammatory:** Aceclofenac inhibits prostaglandin synthesis, thereby reducing pain, swelling and inflammation.
- **Muscle Relaxation:** Tizanidine acts centrally to reduce excessive skeletal muscle tone and muscle spasm.
- **Reduction of Muscle Spasm:** The combination provides relief from pain associated with acute musculoskeletal conditions accompanied by painful muscle spasm.
- **Improved Mobility:** Reduction in pain and muscle spasm may help improve movement and functional mobility.

Indications

Aceflick-TZ Tablets are indicated for the management of:

- Acute painful musculoskeletal conditions associated with muscle spasm
- Acute low back pain with painful muscle spasm
- Neck pain and cervical muscle spasm
- Sprains and strains associated with painful muscle spasm
- Other acute musculoskeletal pain conditions, as prescribed by the physician

Clinical evidence has demonstrated efficacy of the aceclofenac 100 mg + tizanidine 2 mg combination in acute low back pain with muscle spasm.

Dosage and Administration

- **Recommended Dose:** One tablet twice daily or as directed by the physician.

- **Administration:** Swallow the tablet whole with water, preferably after meals to reduce gastrointestinal discomfort.
- **Duration of Use:** Use for the shortest duration necessary and strictly according to the physician's advice.
- Do not exceed the recommended dose.

Precautions

- **Pregnancy & Lactation:** Use during pregnancy or breastfeeding only when specifically prescribed by a healthcare professional.
- **Liver & Kidney Impairment:** Use cautiously in patients with hepatic or renal impairment. Dose adjustment and appropriate monitoring may be required.
- **Gastrointestinal Disorders:** Patients with a history of peptic ulceration or gastrointestinal bleeding should use aceclofenac-containing medicines only after medical assessment.
- **Cardiovascular Conditions:** Use cautiously in patients with hypertension, heart failure or other cardiovascular risk factors.
- **Drowsiness & Dizziness:** Tizanidine may cause drowsiness, dizziness and reduced alertness. Patients should avoid driving or operating machinery if affected.
- **Alcohol:** Avoid alcohol because it may enhance the sedative effects of tizanidine and may increase gastrointestinal risks associated with NSAIDs.

Contraindications

- Known hypersensitivity to aceclofenac, tizanidine, other NSAIDs or any component of the formulation.
- Patients with active or recurrent peptic ulcer disease or gastrointestinal bleeding.
- Patients with severe hepatic impairment.
- Patients with severe renal impairment unless specifically advised and monitored by a physician.
- Concomitant use of tizanidine with potent CYP1A2 inhibitors such as **ciprofloxacin or fluvoxamine** is contraindicated because of the risk of excessive tizanidine exposure and severe hypotension/sedation.

Adverse Reactions

Common:

- Nausea
- Indigestion or stomach discomfort
- Dizziness
- Drowsiness
- Dry mouth
- Weakness or fatigue

Rare but Serious:

- Gastrointestinal bleeding or ulceration
- Severe allergic reactions

- Liver-function abnormalities
- Marked hypotension or bradycardia
- Excessive sedation

Patients should seek medical attention if severe abdominal pain, gastrointestinal bleeding, breathing difficulty, severe dizziness or other serious symptoms occur.

Drug Interactions

- **CYP1A2 inhibitors:** Ciprofloxacin and fluvoxamine can markedly increase tizanidine levels and should not be combined with tizanidine.
- **CNS depressants:** Alcohol, benzodiazepines, opioids and other sedatives may increase drowsiness and CNS depression.
- **Antihypertensive medicines:** Tizanidine may enhance hypotensive effects and blood pressure should be monitored.
- **Other NSAIDs:** Concurrent use with other NSAIDs may increase the risk of gastrointestinal and renal adverse effects.
- **Anticoagulants/antiplatelet agents:** Concomitant use may increase the risk of gastrointestinal bleeding and requires medical supervision.

Storage

- Store in a cool, dry place, preferably **below 30°C**, protected from moisture and excessive heat.
- Keep the tablets in the original packaging until use.
- Keep out of reach of children.

Note

This information is intended as a professional pharmaceutical-literature guideline. **Aceflick-TZ should be used under the supervision of a qualified healthcare professional.** Dosage and duration should be individualized according to the patient's clinical condition and response to treatment.

Manufactured in India for:

Cafoli™
LIFECARE

Cafoli Lifecare Pvt. Ltd.
(An ISO 9001: 2015 Certified Co.)
Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113
Corporate Office :
Plot no. 307, Industrial Area Phase-2,
Panchkula, Haryana -134113
TM: Trademark Applied for