Information For the Use of
Amiodarone (150mg) Injection

Amistyle 150 Injection

AMISTYLE 150 INJECTION

Amiodarone Hydrochloride Injection I.P. 150 mg

COMPOSITION

Each 3 ml ampoule contains:

Amiodarone Hydrochloride I.P. equivalent to Amiodarone 150 mg
Water for Injection I.P. q.s.

Confirm the ampoule volume and exact salt expression against the approved product specification before printing.

PHARMACOLOGICAL ACTION

Amiodarone is a **Class III antiarrhythmic agent** that primarily prolongs cardiac action-potential duration and the refractory period by blocking potassium channels. It also exhibits sodium-channel-blocking, calcium-channel-blocking and non-competitive beta-adrenergic-blocking effects.

These combined actions help stabilize cardiac electrical activity and control serious ventricular and selected supraventricular arrhythmias.

INDICATIONS

Amistyle 150 Injection may be indicated for:

- **Life-threatening recurrent ventricular fibrillation**
- **Haemodynamically unstable ventricular tachycardia**, when clinically appropriate
- Selected serious ventricular arrhythmias that have not responded adequately to other treatments
- Certain acute tachyarrhythmias under specialist supervision, according to applicable guidelines and approved product information

Amiodarone injection is generally used in monitored hospital or emergency-care settings.

DOSAGE AND ADMINISTRATION

Administration must be performed by trained healthcare professionals with continuous cardiac monitoring.

For adult patients with certain serious arrhythmias, commonly referenced IV regimens include:

- **Initial infusion:** 150 mg diluted in 100 ml of 5% dextrose, infused over 10 minutes, for selected ventricular arrhythmias.
- **Subsequent infusion:** May be followed by a continuous infusion according to the applicable protocol and clinical response.

For cardiac arrest due to refractory ventricular fibrillation or pulseless ventricular tachycardia, the resuscitation regimen differs; follow the current advanced life-support protocol and approved product information.

Important administration precautions:

- Do not administer the routine diluted infusion as a rapid IV injection.
- Use an appropriate infusion pump and compatible diluent.
- Monitor ECG, blood pressure and heart rate continuously.
- Monitor electrolytes and correct potassium and magnesium abnormalities.
- The exact dose, dilution, infusion rate and line requirements must follow the approved formulation instructions and hospital protocol.

PRECAUTIONS

- Continuous ECG and blood-pressure monitoring are essential during IV administration.
- Rapid infusion may cause **hypotension, bradycardia or conduction disturbances**.
- Use cautiously in patients with sinus-node dysfunction, AV conduction abnormalities or pre-existing QT prolongation.
- Monitor for hepatic injury and thyroid dysfunction, particularly with ongoing therapy.
- Correct hypokalaemia and hypomagnesaemia before administration where possible.
- Use cautiously in patients with pulmonary disease or a history of serious drug-related pulmonary reactions.
- Review concomitant medicines that prolong the QT interval or slow cardiac conduction.
- Amiodarone can interact with warfarin, digoxin and several other medicines.

CONTRAINDICATIONS

Depending on the approved product information, contraindications may include:

- Hypersensitivity to amiodarone, iodine where relevant to the formulation, or any excipient
- Severe sinus-node dysfunction or significant bradycardia without a functioning pacemaker
- Second- or third-degree AV block without a functioning pacemaker
- Cardiogenic shock or severe hypotension, unless the clinical circumstances and approved emergency indication justify use under specialist care
- Concomitant medicines or conditions specifically contraindicated by the approved product label

---

ADVERSE REACTIONS

Common

- Hypotension
- Bradycardia
- Nausea
- Infusion-site reactions
- Transient conduction disturbances

Serious / Rare

- Severe hypotension or cardiovascular collapse
- Marked bradycardia or AV block
- QT prolongation and, rarely, serious arrhythmias
- Hepatic injury
- Thyroid dysfunction
- Pulmonary toxicity, particularly with prolonged exposure
- Severe hypersensitivity reactions

DRUG INTERACTIONS

Clinically important interactions may occur with:

- **Digoxin:** increased digoxin exposure and risk of bradycardia or conduction disturbances
- **Warfarin:** increased anticoagulant effect and bleeding risk
- **Beta-blockers and certain calcium-channel blockers:** increased risk of bradycardia or AV block
- **Other QT-prolonging medicines:** increased risk of arrhythmias
- **Certain macrolides, azole antifungals and other interacting medicines:** possible changes in arrhythmia risk or amiodarone exposure

- **Certain statins:** increased risk of muscle toxicity with susceptible agents

A complete medication review is recommended before administration.

USE IN PREGNANCY AND LACTATION

Pregnancy: Use only when the potential benefit justifies the potential risk, under specialist supervision. Amiodarone may affect fetal thyroid function and development.

Lactation: Amiodarone and its metabolite may enter breast milk. Breastfeeding decisions require specialist assessment and should follow the approved product information.

STORAGE

- Store according to the approved product label.
- Protect from light.
- Do not freeze unless the approved product instructions specifically permit it.
- Inspect the solution for particles or discoloration before use.
- Keep out of reach of children.

NOTE

Amistyle 150 Injection is intended for **hospital use under qualified medical supervision**. It is not a substitute for immediate electrical cardioversion or defibrillation when these are indicated.

For professional use only.

Manufactured in India for:

Cafoli™
LIFECARE

Cafoli Lifecare Pvt. Ltd.
(An ISO 9001: 2015 Certified Co.)
Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113
Corporate Office :
Plot no. 307, Industrial Area Phase-2,
Panchkula, Haryana -134113
TM: Trademark Applied for