Information For the Use of
Amiodarone (200mg) Tablet

Amistyle 200 Tablet

Amistyle 200 Tablets

Composition:

Each tablet contains:

- **Amiodarone Hydrochloride IP** – 200 mg

Pharmacological Action:

Amistyle 200 Tablets contains **Amiodarone**, a potent antiarrhythmic agent with predominantly Class III activity. It acts on cardiac electrical conduction and helps restore and maintain normal cardiac rhythm.

- **Potassium Channel Blockade:** Prolongs cardiac action potential duration and the effective refractory period.
- **Sodium Channel Blockade:** Slows cardiac conduction and helps suppress abnormal electrical impulses.
- **Calcium Channel Blockade:** Reduces conduction through the atrioventricular (AV) node.
- **Antiadrenergic Action:** Exhibits sympathetic-blocking activity, helping reduce abnormal cardiac excitability.
- **Rhythm Stabilization:** Helps control and prevent recurrent clinically significant cardiac arrhythmias.

Indications:

Amistyle 200 Tablets is indicated for the management of certain serious cardiac arrhythmias, including:

- **Ventricular Tachycardia**
- **Ventricular Fibrillation** in appropriate clinical settings
- **Atrial Fibrillation**
- **Atrial Flutter**
- **Supraventricular Tachyarrhythmias**
- Other clinically significant arrhythmias as determined by the cardiologist

---

Dosage and Administration:

- **Recommended Dose:** Dosage varies according to the type and severity of arrhythmia and should be determined by the physician.
- Amiodarone therapy may require an initial **loading regimen** followed by a lower maintenance dose.
- **Administration:** Swallow the tablet whole with water. It may be taken with or without food, but should preferably be taken consistently in relation to meals.
- Do not discontinue treatment abruptly unless advised by the treating physician.

Precautions:

- Amiodarone requires careful monitoring because of its potential effects on the **thyroid, liver, lungs, eyes, and heart**.
- Baseline and periodic **ECG, liver function, and thyroid function** monitoring may be required.
- Patients should report persistent cough, breathlessness, unexplained weight changes, vision disturbances, or worsening palpitations.
- Use cautiously in patients with **bradycardia, conduction abnormalities, liver disease, or thyroid disorders**.
- Amiodarone can increase sensitivity to sunlight; appropriate sun protection is advisable.
- **Pregnancy and Lactation:** Use only when clearly indicated and under specialist supervision.

Contraindications:

- Known hypersensitivity to **Amiodarone**, iodine, or any component of the formulation.
- Significant **sinus bradycardia** or advanced conduction disorders in the absence of an appropriate pacemaker.
- Certain clinically significant AV or sinoatrial conduction disturbances.
- Concomitant use with certain medicines capable of causing serious cardiac rhythm disturbances, where contraindicated.

Adverse Reactions:

Common:

- Nausea
- Vomiting
- Constipation
- Fatigue
- Photosensitivity
- Bradycardia

Less Common but Serious:

- **Pulmonary toxicity**
- Thyroid dysfunction
- Liver enzyme elevation or hepatic injury
- Corneal microdeposits and visual disturbances
- QT prolongation and proarrhythmia
- Severe skin reactions
- Peripheral neuropathy

Patients should seek medical attention for unexplained breathlessness, persistent cough, fainting, severe weakness, or significant visual changes.

Drug Interactions:

- Concomitant use with other medicines that **prolong the QT interval** may increase the risk of serious arrhythmias.
- Amiodarone can increase the effects or concentrations of **warfarin, digoxin, and certain statins**.
- Concomitant use with other antiarrhythmic agents requires specialist supervision.
- Grapefruit juice may increase amiodarone exposure and should generally be avoided.
- Patients should inform their physician about all prescription medicines, OTC medicines, and supplements being used.

Storage:

- Store in a cool, dry place below **25°C**.
- Protect from moisture and direct sunlight.
- Keep out of reach of children.

Note:

This information is intended as a general prescribing guide. Amistyle 200 Tablets should be used under the supervision of a registered medical practitioner, preferably with appropriate cardiac monitoring. Dosage and duration should be individualized according to the type of arrhythmia, clinical response, and patient's overall condition.

Manufactured in India for:

Cafoli Lifecare Pvt. Ltd.
(An ISO 9001: 2015 Certified Co.)
Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113
Corporate Office :
Plot no. 307, Industrial Area Phase-2,
Panchkula, Haryana -134113
TM: Trademark Applied for