Information For the Use of
Atorvastatin (10mg) + Aspirin (150mg) Capsules

Atorable ASP 10/150 Capsule

ATORABLE ASP 10/150 CAPSULE

Atorvastatin 10 mg + Aspirin 150 mg

COMPOSITION

Each capsule contains:

Atorvastatin Calcium I.P. equivalent to Atorvastatin 10 mg
Aspirin I.P. 150 mg
Excipients q.s.

PHARMACOLOGICAL ACTION

Atorvastatin is an HMG-CoA reductase inhibitor that reduces cholesterol synthesis in the liver, lowers LDL cholesterol and triglycerides, and helps reduce the risk of atherosclerotic cardiovascular events.

Aspirin irreversibly inhibits platelet cyclooxygenase-1 (COX-1), reducing thromboxane A_2 production and platelet aggregation. This helps prevent platelet-rich blood clots.

The combination provides lipid-lowering and antiplatelet effects in patients for whom both treatments are clinically indicated.

INDICATIONS

Atorable ASP 10/150 Capsule may be indicated as part of cardiovascular risk management in appropriately selected patients, including:

- Secondary prevention in patients with established coronary artery disease.
- Prevention of recurrent myocardial infarction.
- Prevention of recurrent ischaemic events in patients for whom antiplatelet therapy is appropriate.
- Management of atherosclerotic cardiovascular disease requiring both lipid-lowering and antiplatelet therapy.
- Selected patients at high cardiovascular risk, when the benefits of both components outweigh the risks.

The combination should be prescribed only when both atorvastatin and aspirin are appropriate for the patient.

DOSAGE AND ADMINISTRATION

Usual adult dosage: One capsule once daily, or as directed by the physician.

- May be taken with or without food.
- Swallow the capsule with water.
- Take it at a consistent time each day.
- Do not discontinue prescribed aspirin used for secondary cardiovascular prevention without medical advice.
- Monitor lipid profile and assess bleeding risk periodically.
- The prescribed regimen should be reviewed if the patient requires a different atorvastatin dose or aspirin dose.

PRECAUTIONS

- Aspirin increases the risk of gastrointestinal and other bleeding.
- Use cautiously in patients with a history of peptic ulcer, gastrointestinal bleeding or bleeding disorders.
- Inform the physician before surgery or dental procedures.
- Avoid unnecessary concomitant use of other NSAIDs or aspirin-containing products.
- Atorvastatin may cause muscle-related adverse effects; report unexplained muscle pain, tenderness or weakness.
- Assess liver function when clinically indicated.
- Review potential interactions that may increase atorvastatin exposure and the risk of myopathy.
- Use cautiously in patients with significant alcohol consumption or a history of liver disease.
- Aspirin may precipitate bronchospasm in susceptible patients with aspirin-sensitive asthma.

CONTRAINDICATIONS

- Hypersensitivity to atorvastatin, aspirin, other salicylates or any formulation component.
- Active pathological bleeding.
- Active gastrointestinal ulceration or significant gastrointestinal bleeding.
- Severe active liver disease or persistent unexplained elevations of liver enzymes.
- Previous serious aspirin/NSAID-induced hypersensitivity, including bronchospasm.
- Pregnancy: atorvastatin should not be used during pregnancy.

ADVERSE REACTIONS

Common

Atorvastatin:

- Muscle pain
- Headache
- Nausea
- Diarrhoea
- Gastrointestinal discomfort

Aspirin:

- Dyspepsia
- Gastric irritation
- Nausea
- Easy bruising
- Increased bleeding tendency

Serious / Less Common

- Gastrointestinal ulceration or bleeding
- Intracranial or other serious bleeding
- Elevated liver enzymes
- Myopathy or myositis
- Rare rhabdomyolysis
- Severe hypersensitivity reactions
- Aspirin-associated bronchospasm

DRUG INTERACTIONS

Atorvastatin interactions:

- Strong CYP3A4 inhibitors, including certain macrolides and azole antifungals, may increase atorvastatin exposure.
- Cyclosporine and some antiviral medicines may increase the risk of muscle toxicity.
- Gemfibrozil, other fibrates and colchicine may increase myopathy risk.
- Large quantities of grapefruit juice may increase atorvastatin exposure.

Aspirin interactions:

- Anticoagulants, such as warfarin and direct oral anticoagulants, may increase bleeding risk.
- Other antiplatelet agents and NSAIDs may increase bleeding risk.
- Corticosteroids and SSRIs/SNRIs may increase gastrointestinal bleeding risk.
- Ibuprofen and certain other NSAIDs may interfere with aspirin's antiplatelet effect depending on timing and use.

USE IN PREGNANCY AND LACTATION

Pregnancy: Atorvastatin is contraindicated during pregnancy. Aspirin use depends on the indication, dose and stage of pregnancy; this fixed-dose combination should not be used during pregnancy without specialist assessment.

Lactation: Breastfeeding is generally not recommended during atorvastatin treatment. Aspirin use during lactation also requires medical assessment.

STORAGE

- Store below 25°C or as specified on the approved product label.
- Protect from moisture and light.
- Keep the capsules in the original packaging.
- Keep out of reach of children.

NOTE

Atorable ASP 10/150 Capsule is intended for patients requiring both lipid-lowering and antiplatelet therapy. The choice of aspirin dose and statin intensity should be individualized according to cardiovascular risk, bleeding risk and clinical guidelines.

For professional use only.

Manufactured in India for:

Cafoli™
LIFECARE

Cafoli Lifecare Pvt. Ltd.
(An ISO 9001: 2015 Certified Co.)
Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113
Corporate Office :
Plot no. 307, Industrial Area Phase-2,
Panchkula, Haryana -134113
TM: Trademark Applied for