Information For the Use of
Atorvastatin (20mg) + Aspirin (150mg) Capsules

Atorable ASP 20/150 Capsule

ATORABLE ASP 20/150 CAPSULE

Atorvastatin 20 mg + Aspirin 150 mg

COMPOSITION

Each hard gelatin capsule contains:

Atorvastatin Calcium I.P. equivalent to Atorvastatin 20 mg
Aspirin I.P. 150 mg
Excipients q.s.

PHARMACOLOGICAL ACTION

Atorvastatin is an HMG-CoA reductase inhibitor. It reduces hepatic cholesterol synthesis and increases hepatic LDL-receptor activity, thereby lowering LDL-C, total cholesterol and triglycerides and producing a modest increase in HDL-C.

Aspirin irreversibly inhibits platelet cyclooxygenase-1 (COX-1), reducing thromboxane A_2 formation and thereby inhibiting platelet aggregation for the lifespan of the platelet.

The combination provides **lipid-lowering and antiplatelet therapy**, commonly used for cardiovascular risk reduction in appropriately selected patients.

INDICATIONS

Atorable ASP 20/150 Capsule may be indicated as part of secondary cardiovascular prevention in appropriate patients, including:

- Prevention of recurrent **myocardial infarction**
- Prevention of recurrent **ischemic stroke or transient ischemic events**, where antiplatelet therapy is appropriate
- Patients with established **coronary artery disease**
- Patients with atherosclerotic cardiovascular disease requiring both statin and antiplatelet therapy
- Prevention of cardiovascular events in appropriately selected high-risk patients

The combination should be used when both components are clinically indicated.

DOSAGE AND ADMINISTRATION

Usual adult dosage:
One capsule once daily, or as directed by the physician.

- May be taken with or without food.
- Preferably take at the same time each day.
- Swallow the capsule whole with water.
- Do not discontinue aspirin abruptly without medical advice when prescribed for secondary cardiovascular prevention.
- Dosage should be individualized according to lipid levels, cardiovascular risk, bleeding risk and clinical response.

Lipid profile and liver function should be monitored according to clinical requirements.

PRECAUTIONS

- Aspirin increases the risk of **gastrointestinal and other bleeding**.
- Use cautiously in patients with a history of peptic ulcer disease or gastrointestinal bleeding.
- Inform the physician before surgery or dental procedures.
- Avoid unnecessary use with other NSAIDs or aspirin-containing medicines.
- Atorvastatin should be used cautiously in patients with liver disease or significant alcohol consumption.
- Report unexplained muscle pain, tenderness or weakness, particularly if associated with fever or malaise.
- Monitor liver enzymes when clinically indicated.
- Certain medicines can significantly increase atorvastatin exposure and the risk of myopathy/rhabdomyolysis.
- Aspirin may precipitate bronchospasm in susceptible patients with aspirin-sensitive asthma.

CONTRAINDICATIONS

- Hypersensitivity to atorvastatin, aspirin, other salicylates or formulation components
- Active pathological bleeding
- Active peptic ulcer disease with clinically significant bleeding
- Severe hepatic disease or persistent unexplained elevation of serum transaminases
- Previous serious aspirin/NSAID-induced asthma, urticaria or hypersensitivity reaction

- Pregnancy, particularly because of atorvastatin; statins are contraindicated during pregnancy
- Breastfeeding is generally not recommended during statin therapy

---

ADVERSE REACTIONS

Common

Atorvastatin:

- Headache
- Gastrointestinal discomfort
- Nausea
- Diarrhea
- Muscle pain

Aspirin:

- Dyspepsia
- Gastric irritation
- Nausea
- Easy bruising
- Increased bleeding tendency

Serious / Rare

- Gastrointestinal ulceration or bleeding
- Intracranial or other serious bleeding
- Hemorrhagic reactions
- Elevated liver enzymes
- Myopathy
- Myositis
- Rare rhabdomyolysis
- Severe hypersensitivity reactions or bronchospasm

DRUG INTERACTIONS

Atorvastatin exposure may increase with strong CYP3A4 inhibitors such as certain macrolide antibiotics and azole antifungals, increasing the risk of myopathy.

Other clinically important interactions include:

- Cyclosporine
- Certain HIV/HCV protease inhibitors
- Gemfibrozil and other fibrates
- Niacin at lipid-lowering doses
- Colchicine
- Grapefruit juice in substantial quantities

Aspirin may increase bleeding risk with:

- Anticoagulants such as warfarin and direct oral anticoagulants
- Other antiplatelet agents
- NSAIDs
- Corticosteroids
- SSRIs/SNRIs and other medicines affecting haemostasis

Ibuprofen and some other NSAIDs may interfere with aspirin's antiplatelet effect when taken concomitantly.

USE IN PREGNANCY AND LACTATION

Pregnancy:
Atorvastatin is contraindicated during pregnancy because cholesterol and other products of the cholesterol biosynthetic pathway are important for fetal development. Aspirin use during pregnancy should be determined according to dose, gestational age and clinical indication.

Lactation:
Breastfeeding is generally not recommended during atorvastatin treatment. Aspirin may pass into breast milk and should be used during lactation only when specifically advised by a physician.

STORAGE

Store below **25°C**, protected from moisture and light.

Keep the capsules in the original container and keep out of reach of children.

NOTE

Atorable ASP 20/150 Capsule should be used under medical supervision when **both lipid-lowering and antiplatelet therapy are clinically indicated**.

Patients should not discontinue antiplatelet therapy prescribed for established cardiovascular disease without consulting their physician.

For professional use only.

Manufactured in India for:

Cafoli™
LIFECARE

Cafoli Lifecare Pvt. Ltd.
(An ISO 9001: 2015 Certified Co.)
Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113
Corporate Office :
Plot no. 307, Industrial Area Phase-2,
Panchkula, Haryana -134113
TM: Trademark Applied for