Atorable ASP 40/75 Capsule

ATORABLE ASP 40/75 CAPSULE

Atorvastatin 40 mg + Aspirin 75 mg

COMPOSITION

Each capsule contains:

Atorvastatin Calcium I.P. equivalent to Atorvastatin 40 mg
Aspirin I.P. 75 mg
Excipients q.s.

PHARMACOLOGICAL ACTION

Atorvastatin is an HMG-CoA reductase inhibitor that reduces hepatic cholesterol synthesis, lowers LDL cholesterol and triglycerides, and helps reduce the risk of atherosclerotic cardiovascular events.

Aspirin irreversibly inhibits platelet cyclooxygenase-1 (COX-1), reducing thromboxane A_2 production and platelet aggregation. This helps prevent the formation of platelet-rich blood clots.

The combination provides lipid-lowering and antiplatelet effects for patients who require both treatments for cardiovascular risk management.

INDICATIONS

Atorable ASP 40/75 Capsule may be indicated for:

- Secondary prevention in patients with established coronary artery disease.
- Prevention of recurrent myocardial infarction.
- Prevention of recurrent ischaemic cardiovascular events when antiplatelet therapy is appropriate.
- Management of atherosclerotic cardiovascular disease requiring intensive lipid-lowering and antiplatelet therapy.
- Selected patients following coronary revascularisation, when both components are indicated.

The combination should be prescribed only when both atorvastatin and aspirin are appropriate for the patient.

DOSAGE AND ADMINISTRATION

Usual adult dosage: One capsule once daily, or as directed by the physician.

- May be taken with or without food.
- Swallow the capsule with water.
- Take it consistently at the same time each day.
- Monitor lipid profile and assess bleeding risk periodically.
- Review treatment if the patient requires a different statin or aspirin dose.
- Do not stop prescribed aspirin used for secondary cardiovascular prevention without medical advice.

PRECAUTIONS

- Aspirin increases the risk of gastrointestinal and other bleeding.
- Use cautiously in patients with a history of peptic ulcer disease, gastrointestinal bleeding or bleeding disorders.
- Inform the physician before surgery or dental procedures.
- Avoid unnecessary concomitant use of other NSAIDs or aspirin-containing products.
- Atorvastatin may cause muscle-related adverse effects. Report unexplained muscle pain, tenderness or weakness.
- Assess liver function when clinically indicated.
- Review medicines that may increase atorvastatin exposure and the risk of myopathy.
- Use cautiously in patients with significant alcohol consumption or a history of liver disease.
- Aspirin may precipitate bronchospasm in patients with aspirin-sensitive asthma.

CONTRAINDICATIONS

- Hypersensitivity to atorvastatin, aspirin, other salicylates or any formulation component.
- Active pathological bleeding.
- Active gastrointestinal ulceration or significant gastrointestinal bleeding.
- Severe active liver disease or persistent unexplained elevations of liver enzymes.
- Previous serious aspirin/NSAID-induced hypersensitivity, including bronchospasm.
- Pregnancy, because atorvastatin should not be used during pregnancy.

ADVERSE REACTIONS

Common

Atorvastatin:

- Muscle pain
- Headache

- Nausea
- Diarrhoea
- Gastrointestinal discomfort

Aspirin:

- Dyspepsia
- Gastric irritation
- Nausea
- Easy bruising
- Increased bleeding tendency

Serious / Less Common

- Gastrointestinal ulceration or bleeding
- Intracranial or other serious bleeding
- Elevated liver enzymes
- Myopathy or myositis
- Rare rhabdomyolysis
- Severe hypersensitivity reactions
- Aspirin-associated bronchospasm

DRUG INTERACTIONS

Atorvastatin interactions:

- Strong CYP3A4 inhibitors, including certain macrolides and azole antifungals, may increase atorvastatin exposure.
- Cyclosporine and some antiviral medicines may increase the risk of muscle toxicity.
- Gemfibrozil, other fibrates and colchicine may increase myopathy risk.
- Large quantities of grapefruit juice may increase atorvastatin exposure.

Aspirin interactions:

- Anticoagulants, including warfarin and direct oral anticoagulants, may increase bleeding risk.
- Other antiplatelet agents and NSAIDs may increase bleeding risk.
- Corticosteroids and SSRIs/SNRIs may increase gastrointestinal bleeding risk.
- Ibuprofen and certain other NSAIDs may interfere with aspirin's antiplatelet effect, depending on timing and use.

USE IN PREGNANCY AND LACTATION

Pregnancy: Atorvastatin is contraindicated during pregnancy. This fixed-dose combination should not be used during pregnancy; aspirin use depends on the indication, dose and stage of pregnancy.

Lactation: Breastfeeding is generally not recommended during atorvastatin treatment. Aspirin use during lactation requires medical assessment.

STORAGE

- Store according to the approved product label, generally below 25°C.
- Protect from moisture and light.
- Keep capsules in the original packaging.
- Keep out of reach of children.

NOTE

Atorable ASP 40/75 Capsule is intended for patients requiring both lipid-lowering and antiplatelet therapy. The choice of statin intensity and aspirin dose should be individualized according to cardiovascular risk, bleeding risk and current clinical guidelines.

For professional use only.

Manufactured in India for:

Cafoli™
LIFECARE

Cafoli Lifecare Pvt. Ltd.
(An ISO 9001: 2015 Certified Co.)
Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113
Corporate Office :
Plot no. 307, Industrial Area Phase-2,
Panchkula, Haryana -134113
TM: Trademark Applied for