Information For the Use of
Atorvastatin (10mg) + Vitamin D3 (1000IU) Tablet

Atorable D 10 Tablet

Atorable D 10 Tablet

Composition:

Each tablet contains:

- **Atorvastatin IP** – 10 mg
- **Vitamin D3 (Cholecalciferol)** – 1000 IU

Pharmacological Action:

Atorable D 10 Tablet combines a lipid-lowering agent with vitamin D supplementation.

- **Lipid-Lowering Action:** Atorvastatin inhibits HMG-CoA reductase, reducing cholesterol synthesis in the liver and increasing hepatic LDL-receptor activity, thereby lowering LDL cholesterol and triglyceride levels.
- **Cardiovascular Risk Reduction:** By reducing LDL cholesterol, atorvastatin helps reduce the risk of cardiovascular events in appropriate patients.
- **Vitamin D Supplementation:** Vitamin D3 supports calcium and phosphorus absorption and helps maintain normal bone and muscle function.
- **Complementary Action:** Provides lipid management along with vitamin D supplementation when both are clinically indicated.

Indications:

Atorable D 10 Tablet is indicated for the management of:

- **Hypercholesterolemia**
- **Mixed Dyslipidemia**
- **Elevated LDL Cholesterol**
- **Prevention of Cardiovascular Events** in patients at appropriate cardiovascular risk
- **Vitamin D Supplementation** in patients requiring supplementation

---

Dosage and Administration:

- **Recommended Dose:** One tablet once daily or as directed by the physician.
- **Administration:** Swallow the tablet whole with water. It may be taken with or without food.
- Take the medicine at the same time each day for better compliance.
- Continue appropriate dietary and lifestyle modifications as advised by the physician.

Precautions:

- Use with caution in patients with **liver disease** or a history of unexplained persistent elevation of liver enzymes.
- Patients should report unexplained muscle pain, tenderness, weakness, or dark-colored urine.
- Avoid excessive intake of vitamin D or calcium supplements unless recommended by the physician.
- Periodic monitoring of lipid profile and liver function may be advised.
- **Pregnancy and Lactation:** Atorvastatin should not be used during pregnancy or breastfeeding.

Contraindications:

- Known hypersensitivity to atorvastatin, vitamin D3, or any component of the formulation.
- **Active liver disease** or persistent unexplained elevation of serum transaminases.
- **Pregnancy and breastfeeding** due to the atorvastatin component.

Adverse Reactions:

Common:

- Headache
- Nausea
- Abdominal discomfort
- Constipation
- Muscle aches

Rare but Serious:

- Significant muscle injury or rhabdomyolysis
- Elevated liver enzymes
- Severe hypersensitivity reactions
- Hypercalcemia with excessive vitamin D intake

Drug Interactions:

- Strong CYP3A4 inhibitors, including certain macrolide antibiotics and azole antifungals, may increase atorvastatin exposure and the risk of muscle toxicity.
- Concomitant use with fibrates may increase the risk of myopathy.
- Excessive consumption of grapefruit or grapefruit juice may increase atorvastatin exposure.
- Certain medicines affecting calcium or vitamin D metabolism may alter the effects of vitamin D3.

- Inform the physician about all prescription medicines, OTC medicines, and supplements being used.

Storage:

- Store in a cool, dry place below **25°C**.
- Protect from moisture and direct sunlight.
- Keep out of reach of children.

Note:

This information is intended as a general prescribing guide. Atorable D 10 Tablet should be used under the supervision of a registered medical practitioner. The dosage and duration of treatment should be individualized according to the patient's lipid profile, cardiovascular risk, vitamin D requirements, and clinical condition.

Manufactured in India for:

Cafoli™
LIFECARE

Cafoli Lifecare Pvt. Ltd.
(An ISO 9001: 2015 Certified Co.)
Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113
Corporate Office :
Plot no. 307, Industrial Area Phase-2,
Panchkula, Haryana -134113
TM: Trademark Applied for