Information For the Use of
Azathioprine 50mg Tablet

Azathipair 50 Tablet

Brand Name: Azathipair 50 Tablet
Dosage Form: Tablet
Composition:
Each tablet contains:
• Azathioprine IP 50 mg

Description:
Azathipair 50 Tablet contains Azathioprine, an immunosuppressive agent belonging to the purine analogue class. It is indicated for the prevention of organ transplant rejection and for the management of various autoimmune and inflammatory disorders. Azathioprine suppresses the proliferation of lymphocytes by inhibiting purine synthesis, thereby reducing immune-mediated tissue damage and controlling disease activity.

Indications:
Indicated for the treatment and management of:
• Prevention of organ transplant rejection
• Rheumatoid arthritis
• Systemic lupus erythematosus (SLE)
• Inflammatory bowel disease (Crohn's disease and ulcerative colitis)
• Autoimmune hepatitis
• Nephrotic syndrome
• Other autoimmune disorders as determined by the physician
It helps suppress abnormal immune responses, reduce inflammation, and maintain disease remission.

Dosage & Administration:
Dosage should be individualized and administered strictly as directed by the physician.
The tablet should be swallowed whole with water and may be taken after meals to reduce gastrointestinal discomfort.
Regular monitoring of complete blood count (CBC), liver function, and renal function is recommended during therapy.
Do not discontinue treatment without medical advice.

Pharmacological Action:

- **Azathioprine is a prodrug that is converted to 6-mercaptopurine and subsequently to active metabolites that inhibit de novo purine synthesis.**
- **This suppresses the proliferation of T and B lymphocytes, reducing immune cell activity and antibody production.**
- **The immunosuppressive action helps prevent organ rejection and controls autoimmune inflammatory diseases.**

Side Effects:
Common side effects include nausea, vomiting, loss of appetite, abdominal discomfort, diarrhea, and mild liver enzyme elevation.
Serious adverse effects may include bone marrow suppression, leukopenia, thrombocytopenia, anemia, increased susceptibility to infections, pancreatitis, hepatotoxicity, and hypersensitivity

reactions. Long-term therapy may increase the risk of certain malignancies, particularly lymphomas and skin cancers.

Precautions & Warnings:

• Use only under the supervision of a qualified physician experienced in immunosuppressive therapy.
• Regular monitoring of complete blood count, liver function, and renal function is essential.
• Screen for TPMT (Thiopurine Methyltransferase) and/or NUDT15 deficiency where appropriate before initiating therapy, as deficiency may increase the risk of severe myelosuppression.
• Avoid live vaccines during treatment.
• Patients should minimize excessive exposure to sunlight and use appropriate sun protection due to an increased risk of skin cancer.
• Use with caution in patients with hepatic or renal impairment.
• Pregnancy and lactation use should be based on a careful benefit-risk assessment by the physician.

Drug Interactions:

May interact with allopurinol, febuxostat, ACE inhibitors, warfarin, ribavirin, mesalamine, sulfasalazine, co-trimoxazole, and other immunosuppressive or myelosuppressive agents. Dose adjustment and close clinical monitoring may be required during concomitant therapy.

Storage:

Store below 25°C in a cool, dry place.
Protect from light and moisture.
Keep out of reach of children.

Manufactured in India for:

Cafoli™
LIFECARE

Cafoli Lifecare Pvt. Ltd.
(An ISO 9001: 2015 Certified Co.)
Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113
Corporate Office :
Plot no. 307, Industrial Area Phase-2,
Panchkula, Haryana -134113
TM: Trademark Applied for