Information For the Use of
Betamethasone (4mg/ml) Injection

Betapair Injection

BETAPAIR INJECTION

Betamethasone Injection 4 mg/ml

COMPOSITION

Each ml contains:

Betamethasone 4 mg
In a suitable sterile vehicle q.s.

Confirm whether the formulation contains betamethasone sodium phosphate or another betamethasone salt, and verify the equivalent strength against the approved product label before printing.

PHARMACOLOGICAL ACTION

Betamethasone is a potent synthetic glucocorticoid with anti-inflammatory, antiallergic and immunosuppressive properties. It acts by binding to intracellular glucocorticoid receptors and modifying gene expression, thereby reducing the production of inflammatory mediators and suppressing immune responses.

It helps reduce inflammation, swelling and allergic manifestations in appropriately selected conditions.

INDICATIONS

Betapair Injection may be indicated for:

- Severe allergic and inflammatory conditions requiring systemic corticosteroid therapy.
- Acute exacerbations of selected inflammatory disorders.
- Severe dermatological conditions when systemic corticosteroid treatment is appropriate.
- Selected rheumatic and musculoskeletal inflammatory conditions.
- Certain autoimmune or connective-tissue disorders as part of specialist-directed treatment.
- Other corticosteroid-responsive conditions when oral therapy is unsuitable or parenteral treatment is clinically indicated.

The approved indications depend on the specific salt, formulation and permitted route of administration.

DOSAGE AND ADMINISTRATION

The dose must be individualized according to the condition, severity, patient response and formulation.

- Administer only by the route specified in the approved product information.
- The dose and duration should be limited to the minimum required to control the condition.
- Intravenous administration, if permitted for the specific formulation, must follow the approved instructions.
- **Do not administer intrathecally or epidurally.**
- Do not inject into a joint or soft tissue unless the particular formulation is approved for that route.
- Long-term or repeated treatment should not be stopped abruptly without medical advice.

PRECAUTIONS

- Use cautiously in patients with diabetes, hypertension, heart failure, osteoporosis or peptic ulcer disease.
- Systemic corticosteroids may increase susceptibility to infections and mask their signs.
- Use cautiously in patients with active or latent tuberculosis and other significant infections.
- Monitor blood glucose, blood pressure and electrolytes when clinically indicated.
- Prolonged use may suppress adrenal function and affect growth in children.
- Avoid live vaccines during immunosuppressive corticosteroid treatment unless specifically advised by a specialist.
- Use cautiously in patients with glaucoma, psychiatric disorders or a history of steroid-related adverse effects.
- Pregnancy and lactation require an individualized benefit–risk assessment.

CONTRAINDICATIONS

- Hypersensitivity to betamethasone or any component of the formulation.
- Systemic fungal infection, unless specifically indicated as part of specialist-directed treatment.
- Administration by a route not approved for the specific formulation.
- Other contraindications specified in the approved product information.

ADVERSE REACTIONS

Common

- Increased appetite
- Indigestion
- Insomnia
- Mood changes
- Increased blood glucose
- Fluid retention
- Increased blood pressure

Serious / Less Common

- Adrenal suppression
- Increased susceptibility to infection
- Severe psychiatric reactions
- Gastrointestinal ulceration or bleeding
- Osteoporosis with prolonged use
- Hypokalaemia
- Severe hypersensitivity reactions
- Injection-site reactions

The risk of adverse effects generally increases with higher doses and prolonged systemic exposure.

DRUG INTERACTIONS

- **Antidiabetic medicines:** Betamethasone may increase blood glucose and require treatment adjustment.
- **NSAIDs:** Increased risk of gastrointestinal irritation, ulceration or bleeding.
- **Anticoagulants:** Anticoagulant response may change and require monitoring.
- **Rifampicin, phenytoin and carbamazepine:** May reduce corticosteroid exposure.
- **Strong CYP3A4 inhibitors**, including certain azole antifungals and macrolides, may increase corticosteroid exposure.
- **Live vaccines:** Increased risk of vaccine-related complications during immunosuppressive therapy.

USE IN PREGNANCY AND LACTATION

Use during pregnancy only when clinically indicated after assessment of potential benefits and risks.

Betamethasone may enter breast milk. Repeated or high-dose treatment during lactation should be assessed by a healthcare professional.

STORAGE

- Store according to the approved product label.
- Protect from excessive heat and light.
- Do not freeze unless specifically permitted by the manufacturer.
- Inspect the solution before use; do not use if it shows unexpected discoloration or visible particles.
- Keep out of reach of children.

NOTE

Betapair Injection should be administered by a qualified healthcare professional. The **exact betamethasone salt, concentration, approved route and dosing instructions must be verified against the registered Indian product information before final printing**.

For professional use only.

Manufactured in India for:

Cafoli™ LIFECARE

Cafoli Lifecare Pvt. Ltd.
(An ISO 9001: 2015 Certified Co.)
Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113
Corporate Office :
Plot no. 307, Industrial Area Phase-2,
Panchkula, Haryana -134113
TM: Trademark Applied for