Information For the Use of
Glycopyrrolate (9mcg) + Formoterol (4.80mcg) + Budesonide (160mcg) Inhaler

Buditerm Plus Inhaler

Brand Name: Buditerm Plus Inhaler
Dosage Form: Inhaler
Composition:
Each metered dose contains:
• Glycopyrrolate IP 9 mcg
• Formoterol Fumarate IP equivalent to Formoterol 4.80 mcg
• Budesonide IP 160 mcg

Description:
Buditerm Plus Inhaler is a triple combination inhalation therapy containing Glycopyrrolate, Formoterol, and Budesonide. Glycopyrrolate is a long-acting muscarinic antagonist (LAMA), Formoterol is a long-acting β_2-adrenergic agonist (LABA), and Budesonide is an inhaled corticosteroid (ICS). This combination provides bronchodilation, reduces airway inflammation, improves lung function, and helps control respiratory symptoms in chronic airway diseases requiring triple therapy.

Indications:
Indicated for the maintenance treatment of:
• Chronic obstructive pulmonary disease (COPD)
• Chronic bronchitis associated with airflow limitation
• Emphysema-associated respiratory symptoms
• Patients requiring inhaled corticosteroid, long-acting bronchodilator, and anticholinergic combination therapy
It helps improve airflow, reduce breathlessness, control airway inflammation, and decrease the frequency of respiratory exacerbations.

Dosage & Administration:
Dosage should be as directed by the physician.
Administer by oral inhalation only using the prescribed inhaler device.
Shake the inhaler well before use (if applicable).
Inhale the medication correctly using proper inhalation technique.
Rinse the mouth with water after each use to reduce the risk of oral fungal infection.
Do not exceed the prescribed dosage.

Pharmacological Action:

- **Glycopyrrolate is a long-acting muscarinic antagonist that blocks M3 receptors in airway smooth muscle, reducing bronchoconstriction and producing prolonged bronchodilation.**
- **Formoterol is a long-acting selective β_2-adrenergic receptor agonist that relaxes bronchial smooth muscle, resulting in rapid and sustained bronchodilation.**
- **Budesonide is an inhaled corticosteroid that suppresses airway inflammation by reducing inflammatory mediator release, airway edema, and hyperresponsiveness.**

Together, these three components provide complementary effects by improving airway patency, reducing inflammation, and enhancing respiratory function.

Side Effects:
Common side effects include throat irritation, cough, hoarseness, dry mouth, headache, tremor,

palpitations, nausea, and oral candidiasis.
Rarely, paradoxical bronchospasm, hypersensitivity reactions, increased heart rate, or systemic corticosteroid-related effects may occur.

Precautions & Warnings:
• For maintenance therapy only; not intended for immediate relief of acute bronchospasm.
• Use with caution in patients with cardiovascular disorders, glaucoma, urinary retention, diabetes mellitus, hyperthyroidism, or active infections.
• Rinse mouth after inhalation to reduce the risk of oral candidiasis.
• Do not discontinue therapy abruptly without medical advice.
• Regular monitoring of lung function and clinical response is recommended during long-term treatment.

Drug Interactions:
May interact with beta-blockers, other sympathomimetic agents, diuretics, xanthine derivatives, corticosteroids, anticholinergic medications, and strong CYP3A4 inhibitors.
Concurrent use with other bronchodilators should be under medical supervision.

Storage:
Store below 25°C in a cool, dry place.
Protect from direct sunlight and excessive heat.
Do not puncture or expose the inhaler canister to fire, even when empty.
Keep out of reach of children.

Manufactured in India for:

Cafoli™
LIFECARE

Cafoli Lifecare Pvt. Ltd.
(An ISO 9001: 2015 Certified Co.)
Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113
Corporate Office :
Plot no. 307, Industrial Area Phase-2,
Panchkula, Haryana -134113
TM: Trademark Applied for