Information For the Use of
Hyoscine butylbromide (10mg) + Paracetamol (325mg) Tablets

Butyhype P Tablet

BUTYHYPE P TABLET

Composition

Each film-coated tablet contains:

Hyoscine Butylbromide I.P. 10 mg
Paracetamol I.P. 325 mg
Excipients … q.s.

Pharmacological Action

Hyoscine Butylbromide is a peripherally acting antispasmodic with anticholinergic activity. It relaxes smooth muscle and helps relieve painful spasms in the gastrointestinal, biliary, urinary and genitourinary tracts.

Paracetamol is an analgesic and antipyretic that provides relief from mild to moderate pain and reduces fever.

The combination provides **antispasmodic and analgesic action**, making it useful for painful conditions associated with smooth-muscle spasm.

Indications

Butyhype P Tablet is indicated for the symptomatic relief of:

- Spasmodic abdominal pain and cramps
- Gastrointestinal tract spasm
- Painful intestinal cramps
- Functional abdominal spasm
- Spasmodic pain associated with the biliary and urinary tracts
- Dysmenorrhoea and other painful spasmodic conditions

The combination is specifically described for spasmodic pain involving the stomach/intestine and functional disorders of the biliary, urinary and female genital tracts.

Dosage and Administration

Adults: 1–2 tablets, up to **3 times daily**, or as directed by the physician.

Maximum: Do not exceed **6 tablets in 24 hours** unless specifically directed by a physician.

Swallow the tablet whole with an adequate amount of water. Do not chew.

Treatment should generally be limited to the shortest duration necessary. Persistent or unexplained abdominal pain should be medically evaluated rather than treated continuously with the product.

Precautions

- Use cautiously in patients with tachycardia, cardiac disease or hyperthyroidism.
- Use with caution in patients prone to intestinal or urinary tract obstruction.
- Patients with severe or unexplained abdominal pain, fever, vomiting, abdominal tenderness, altered bowel movements, fainting or blood in stools should seek immediate medical evaluation.
- Avoid exceeding the recommended paracetamol dose because excessive paracetamol may cause serious liver injury.
- Patients should avoid taking other medicines containing **paracetamol/acetaminophen** concurrently unless advised by a healthcare professional.
- Use cautiously in patients with hepatic or renal impairment.
- Prolonged or excessive alcohol consumption may increase the risk of paracetamol-related liver toxicity.
- Anticholinergic effects may cause dry mouth, constipation, blurred vision or difficulty in urination.

Contraindications

Butyhype P Tablet is contraindicated in patients with:

- Hypersensitivity to hyoscine butylbromide, paracetamol or any component of the formulation
- Myasthenia gravis
- Mechanical stenosis/obstruction of the gastrointestinal tract
- Paralytic or obstructive ileus
- Megacolon
- Narrow-angle glaucoma
- Severe hepatic impairment, where applicable to the paracetamol component

Adverse Reactions

Common/possible:

- Dry mouth
- Constipation
- Nausea
- Dizziness
- Headache
- Blurred vision
- Difficulty in urination
- Increased heart rate

Rare/serious:

- Hypersensitivity reactions
- Skin rash or urticaria
- Severe allergic reactions/anaphylaxis
- Acute increase in intraocular pressure in susceptible patients
- Liver injury following paracetamol overdose or excessive use

Drug Interactions

Hyoscine butylbromide may have enhanced anticholinergic effects when used with:

- Tricyclic or tetracyclic antidepressants
- Antihistamines
- Antipsychotics
- Quinidine or disopyramide
- Amantadine
- Other anticholinergic medicines such as atropine, ipratropium or tiotropium

Concomitant use with **metoclopramide** may reduce the gastrointestinal effects of both medicines. Beta-adrenergic medicines may enhance tachycardic effects.

Paracetamol may interact with medicines such as warfarin and other hepatotoxic medicines, particularly with prolonged use or at higher doses. Patients should inform their healthcare professional about all medicines containing paracetamol.

Pregnancy and Lactation

Use during pregnancy or lactation should be based on medical advice after assessment of the potential benefits and risks.

Storage

Store below **25°C**, in a cool, dry place and protected from excessive heat and moisture.

Keep the medicine in its original packaging and out of reach of children.

Note

For use under medical supervision.

The recommended dose should not be exceeded. The total daily intake of paracetamol from all medicines should be considered to avoid accidental overdose. Persistent, severe or unexplained abdominal pain requires appropriate medical evaluation.

Manufactured in India for:

Cafoli Lifecare Pvt. Ltd.
(An ISO 9001: 2015 Certified Co.)
Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113
Corporate Office :
Plot no. 307, Industrial Area Phase-2,
Panchkula, Haryana -134113
TM: Trademark Applied for