Information For the Use of
Clopidogrel 75mg + Aspirin 150mg Tablet

# Clopiside ASP 150 Tablet

**Brand Name: Clopiside ASP 150 Tablet**
**Dosage Form: Tablet**
**Composition:**
**Each film-coated tablet contains:**
**• Clopidogrel Bisulfate IP equivalent to Clopidogrel 75 mg**
**• Aspirin IP 150 mg**

---

**Description:**
**Clopiside ASP 150 Tablet is a combination of Clopidogrel and Aspirin, both of which possess antiplatelet activity. It is indicated for the prevention and management of thrombotic cardiovascular and cerebrovascular events in high-risk patients. It helps in inhibiting platelet aggregation and reducing the formation of blood clots, thereby improving cardiovascular protection.**

---

**Indications:**
**Indicated for the prevention and treatment of:**
**• Acute coronary syndrome (ACS)**
**• Myocardial infarction**
**• Ischemic stroke**
**• Peripheral arterial disease**
**• Prevention of recurrent thrombotic events**
**It helps in reducing the risk of heart attack, stroke, and vascular complications.**

---

**Dosage & Administration:**
**Dosage should be as directed by the physician.**
**The tablet should be swallowed whole with water, preferably after meals to reduce gastric irritation.**
**Patients should be monitored periodically for bleeding complications.**

---

**Pharmacological Action:**

- **Clopidogrel irreversibly inhibits the platelet P2Y12 ADP receptor, thereby preventing platelet activation and aggregation.**
- **Aspirin inhibits cyclooxygenase-1 (COX-1), reducing thromboxane A2 production and platelet aggregation.**

**Together, these mechanisms provide synergistic dual antiplatelet action for effective prevention of thrombotic events.**

---

**Side Effects:**
**Common side effects include dyspepsia, abdominal discomfort, nausea, bruising, and increased bleeding tendency.**
**Rarely, gastrointestinal bleeding, hypersensitivity reactions, or thrombocytopenia may occur.**

---

**Precautions & Warnings:**
**• Use with caution in patients with peptic ulcer disease or bleeding disorders**
**• Monitor for signs of gastrointestinal or internal bleeding**
**• Avoid unnecessary concomitant use with other NSAIDs or anticoagulants**
**• Use cautiously in hepatic or renal impairment**

**• Discontinue prior to surgery only under medical advice**

---

**Drug Interactions:**
**May interact with anticoagulants, thrombolytics, NSAIDs, proton pump inhibitors, and selective serotonin reuptake inhibitors (SSRIs).**
**Concurrent use with other antiplatelet agents may increase bleeding risk.**

---

**Storage:**
**Store below 25°C in a cool, dry place.**
**Protect from light and moisture.**
**Keep out of reach of children.**

Manufactured in India for:

Cafoli™
LIFECARE

**Cafoli Lifecare Pvt. Ltd.**
**(An ISO 9001: 2015 Certified Co.)**
Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113
TM: Trademark Applied for