Information For the Use of
Clopidogrel (75mg) + Aspirin (75mg) Tablet

Clopiside ASP 75 Tablet

Brand Name: Clopiside ASP 75 Tablet
Dosage Form: Tablet
Composition:
Each film-coated tablet contains:
• Clopidogrel Bisulfate IP equivalent to Clopidogrel 75 mg
• Aspirin IP 75 mg

Description:
Clopiside ASP 75 Tablet is a fixed-dose combination of Clopidogrel, an antiplatelet agent, and Aspirin, a salicylate with antiplatelet activity. It is indicated for the prevention of atherothrombotic events in patients at cardiovascular risk. It helps in inhibiting platelet aggregation and reducing the risk of clot formation associated with cardiovascular and cerebrovascular disorders.

Indications:
Indicated for the prevention and management of:
• Acute coronary syndrome (ACS)
• Myocardial infarction
• Ischemic stroke
• Peripheral arterial disease
• Prevention of thrombotic cardiovascular events
It helps in reducing the risk of heart attack, stroke, and other thromboembolic complications.

Dosage & Administration:
Dosage should be as directed by the physician.
The tablet should be swallowed whole with water, preferably after meals to minimize gastric irritation.
Regular monitoring may be required in patients at risk of bleeding.

Pharmacological Action:

- **Clopidogrel selectively inhibits ADP-induced platelet aggregation by irreversibly blocking the P2Y12 receptor on platelets.**
- **Aspirin inhibits cyclooxygenase-1 (COX-1), thereby reducing thromboxane A2 synthesis and platelet activation.**

Together, these actions provide dual antiplatelet activity for enhanced protection against thrombotic events.

Side Effects:
Common side effects include gastric irritation, abdominal pain, nausea, bruising, and bleeding tendency.
Rarely, gastrointestinal bleeding, allergic reactions, or thrombocytopenia may occur.

Precautions & Warnings:
• Use with caution in patients with bleeding disorders or peptic ulcer disease
• Monitor for signs of unusual bleeding or bruising
• Avoid use before surgical procedures unless advised by the physician

• Use cautiously in hepatic or renal impairment
• Not recommended in patients hypersensitive to aspirin or clopidogrel

---

Drug Interactions:
May interact with anticoagulants, NSAIDs, proton pump inhibitors, and thrombolytic agents.
Concurrent use with other antiplatelet drugs may increase bleeding risk.

Storage:
Store below 25°C in a cool, dry place.
Protect from moisture and light.
Keep out of reach of children.

Manufactured in India for:

Cafoli™
LIFECARE

Cafoli Lifecare Pvt. Ltd.
(An ISO 9001: 2015 Certified Co.)
Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113
TM: Trademark Applied for