Information For the Use of
Levosalbutamol (50mcg)_CFC Free Inhaler

Critsal 50 Inhaler

Critsal 50 Inhaler

Composition:

Each metered dose contains:

- **Levosalbutamol** – 50 mcg
- **CFC Free Propellant**

Pharmacological Action:

Levosalbutamol is a **selective short-acting β_2-adrenergic receptor agonist (SABA)** that provides rapid bronchodilation.

- **Bronchodilation:** Relaxes bronchial smooth muscle, resulting in widening of the airways.
- **Rapid Relief:** Provides prompt relief from bronchospasm and associated breathing difficulty.
- **Improved Airflow:** Reduces airway resistance and improves airflow through the bronchial passages.
- **Selective β_2 Activity:** Primarily acts on β_2 receptors in bronchial smooth muscle.

Indications:

Critsal 50 Inhaler is indicated for the relief and prevention of bronchospasm associated with:

- **Bronchial Asthma**
- **Chronic Obstructive Pulmonary Disease (COPD)**
- **Reversible Airway Obstruction**
- **Exercise-Induced Bronchospasm**, when prescribed

Dosage and Administration:

- **Recommended Dose:** Usually **1–2 inhalations as required**, or as directed by the physician.
- Shake the inhaler well before each use.
- Remove the cap and breathe out fully, away from the mouthpiece.
- Place the mouthpiece between the lips and begin breathing in slowly and deeply while pressing the canister once.
- Hold the breath for approximately **10 seconds**, or as comfortably possible.
- If another inhalation is required, wait approximately **1 minute** before repeating.
- Use a **spacer device** when recommended.

- Do not exceed the prescribed frequency of use.

Precautions:

- Increasing frequency of use or reduced response may indicate **worsening asthma or respiratory disease** and requires medical reassessment.
- Use cautiously in patients with **cardiovascular disease, hypertension, hyperthyroidism, diabetes mellitus, or seizure disorders**.
- May cause tremor, palpitations, or transient increases in heart rate.
- Excessive use may result in **hypokalemia** and other systemic β_2-agonist effects.
- If acute symptoms are not adequately relieved by the prescribed dose, seek immediate medical attention.
- **Pregnancy and Lactation:** Use only when clearly indicated and under medical supervision.

Contraindications:

- Known hypersensitivity to **levosalbutamol, salbutamol, or any component of the formulation**.
- Patients with significant cardiovascular disease, hyperthyroidism, diabetes, or hypokalemia require careful medical supervision rather than routine avoidance.

Adverse Reactions:

Common:

- Tremor
- Headache
- Nervousness
- Palpitations
- Increased heart rate
- Muscle cramps
- Throat irritation

Less Common but Serious:

- Paradoxical bronchospasm
- Significant tachycardia or cardiac arrhythmia
- Hypokalemia
- Hyperglycemia
- Hypersensitivity reactions

If breathing suddenly becomes worse immediately after inhalation, discontinue use and seek urgent medical attention.

Drug Interactions:

- Other **β_2-agonists** may increase cardiovascular and metabolic adverse effects.
- **Non-selective beta-blockers** may reduce the bronchodilator effect and may precipitate bronchospasm.
- **Diuretics** may increase the risk of hypokalemia.
- **MAO inhibitors and tricyclic antidepressants** may enhance cardiovascular effects.
- **Digoxin:** β_2-agonists may contribute to hypokalemia and increase arrhythmia risk in susceptible patients.
- Other sympathomimetic medicines may increase cardiovascular adverse effects.

Storage:

- Store according to the manufacturer's specified conditions, generally below **25°C**.
- Protect from direct sunlight and excessive heat.
- Do not puncture, crush, or incinerate the canister, even when empty.
- Keep the inhaler out of reach of children.
- Keep the mouthpiece clean and replace the protective cap after use.

Note:

This information is intended as a general prescribing guide. **Critsal 50 Inhaler should be used under the supervision of a registered medical practitioner and with correct inhaler technique.** Increasing reliance on a short-acting bronchodilator may indicate deterioration of asthma or COPD and should prompt medical reassessment.

Important: The product's final IP literature should state the exact **metered dose/delivered dose** as specified in the approved product dossier and label.

Manufactured in India for:

Cafoli™ LIFECARE

Cafoli Lifecare Pvt. Ltd.
(An ISO 9001: 2015 Certified Co.)
Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113
Corporate Office :
Plot no. 307, Industrial Area Phase-2,

Panchkula, Haryana -134113
TM: Trademark Applied for