Information For the Use of
Cyclosporine (100mg) Capsules

Cyclonip 100 Capsule

Cyclonip 100 Capsule

Composition:

Each capsule contains:

- **Cyclosporine IP** – 100 mg

Pharmacological Action:

Cyclosporine is a **potent immunosuppressive agent** that selectively suppresses T-lymphocyte-mediated immune responses.

- **Calcineurin Inhibition:** Inhibits calcineurin activity, reducing activation of T-lymphocytes.
- **Reduced Cytokine Production:** Suppresses interleukin-2 and other cytokines involved in immune activation.
- **Immunosuppressive Action:** Reduces immune-mediated tissue injury and helps prevent rejection of transplanted organs.
- **Anti-inflammatory/Immunomodulatory Action:** Its immunosuppressive effects are also utilized in selected autoimmune and inflammatory disorders.

Indications:

Cyclonip 100 Capsule is indicated for:

- **Prevention of Organ Rejection following Kidney, Liver, and Heart Transplantation**
- **Rheumatoid Arthritis** in selected patients inadequately responsive to conventional therapy
- **Severe Plaque Psoriasis** in appropriately selected patients
- Other autoimmune or immune-mediated conditions as determined by a specialist

---

Dosage and Administration:

- **Recommended Dose:** The dose is **individualized according to the indication, body weight, clinical response, and blood cyclosporine concentration**.
- Take consistently at the **same time each day** and in relation to meals.
- Swallow the capsule whole with water.
- Do not crush or chew the capsule.
- Do not switch between different cyclosporine formulations or brands without medical supervision.

- Regular monitoring of **blood cyclosporine levels, renal function, blood pressure, liver function, and electrolytes** is required.
- Do not stop treatment abruptly unless specifically advised by the treating specialist.

Precautions:

- Cyclosporine has a **narrow therapeutic range** and requires specialist supervision.
- Regular monitoring of **serum creatinine and renal function** is essential because nephrotoxicity can occur.
- Monitor **blood pressure, potassium, magnesium, liver function, and lipid levels** during prolonged treatment.
- Increased susceptibility to **bacterial, viral, fungal, and opportunistic infections** may occur.
- Long-term immunosuppression increases the risk of **malignancies, particularly skin cancers and lymphoproliferative disorders**.
- Avoid excessive exposure to sunlight and use appropriate sun protection.
- Avoid **grapefruit and grapefruit juice**, which may alter cyclosporine concentrations.
- Live vaccines may produce inadequate immune responses or may pose a risk during immunosuppressive therapy.
- **Pregnancy and Lactation:** Use only when clearly indicated and under specialist supervision.

Contraindications:

- Known hypersensitivity to **cyclosporine or any component of the formulation**.
- Concomitant use with medicines known to cause serious interactions with cyclosporine should be avoided.
- Severe uncontrolled hypertension or significant renal impairment may preclude use depending on the indication and specialist assessment.
- Cyclosporine should not be used for conditions where the risks of immunosuppression outweigh the expected clinical benefit.

Adverse Reactions:

Common:

- Headache
- Tremor
- Increased hair growth
- Gum enlargement
- Hypertension
- Nausea
- Diarrhea
- Increased serum creatinine
- Increased potassium levels

Less Common but Serious:

- **Nephrotoxicity**
- Hepatotoxicity
- Serious infections
- Severe hypertension
- Hyperkalemia
- Neurotoxicity, including seizures or confusion
- Increased risk of malignancies
- Serious hypersensitivity reactions

Seek immediate medical attention for reduced urine output, severe headache, confusion, seizures, fever, unusual infections, or significant swelling.

Drug Interactions:

Cyclosporine has numerous clinically significant interactions.

- **Macrolide antibiotics** such as erythromycin and clarithromycin may increase cyclosporine concentrations.
- **Azole antifungals** may significantly increase cyclosporine exposure.
- **Rifampicin and certain anticonvulsants** may decrease cyclosporine concentrations and reduce therapeutic effectiveness.
- **NSAIDs and other nephrotoxic medicines** may increase the risk of kidney injury.
- **Statins** may have an increased risk of myopathy/rhabdomyolysis when combined with cyclosporine; selection and dosing require medical supervision.
- **Potassium supplements and potassium-sparing diuretics** may increase the risk of hyperkalemia.
- **Grapefruit/grapefruit juice** may increase cyclosporine concentrations.
- Live vaccines may be inappropriate during treatment.
- Always inform the physician about all prescription medicines, OTC products, herbal preparations, and supplements.

Storage:

- Store in a cool, dry place below **25°C**, unless otherwise specified by the manufacturer.
- Protect from moisture and direct sunlight.
- Keep capsules in the original packaging until use.
- Keep out of reach of children.

Note:

This information is intended as a general prescribing guide. **Cyclonip 100 Capsule should be prescribed and monitored by a specialist experienced in immunosuppressive therapy.** Cyclosporine dosing must be individualized and may require therapeutic drug monitoring.

Important: Cyclosporine has a **narrow therapeutic index**. Patients should not change the dose, brand/formulation, or dosing schedule without medical advice. Regular monitoring of **renal function, blood pressure, liver function, electrolytes, and cyclosporine blood levels** is essential.



Manufactured in India for:

Cafoli™ LIFECARE

Cafoli Lifecare Pvt. Ltd.
(An ISO 9001: 2015 Certified Co.)
Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113
Corporate Office :
Plot no. 307, Industrial Area Phase-2,
Panchkula, Haryana -134113
TM: Trademark Applied for