Information For the Use of
Dabigatran Etexilate 110mg Capsules

Dabiflick 110 Capsule

DABIFLICK 110 CAPSULES

Dabigatran Etexilate 110 mg

Composition

Each hard capsule contains:

- **Dabigatran Etexilate – 110 mg**

Pharmacological Action

Dabigatran etexilate is an orally administered **direct thrombin inhibitor (DTI)**. It is converted to active dabigatran, which directly inhibits thrombin and thereby prevents the conversion of fibrinogen to fibrin.

It functions by:

- **Direct Thrombin Inhibition:** Directly inhibits thrombin (Factor IIa), an important enzyme in the coagulation cascade.
- **Anticoagulant Action:** Reduces formation of fibrin and development of blood clots.
- **Prevention of Thrombus Formation:** Helps prevent pathological clot formation in patients at increased risk of thromboembolic events.
- **Predictable Anticoagulation:** Provides oral anticoagulant activity without routine INR monitoring as required with vitamin K antagonists, although appropriate clinical and renal monitoring remains important. Medicines

Indications

Dabiflick 110 Capsules may be indicated for:

- **Prevention of stroke and systemic embolism in adult patients with non-valvular atrial fibrillation (NVAF)** and appropriate thromboembolic risk.
- **Treatment of deep vein thrombosis (DVT) and pulmonary embolism (PE)** and prevention of recurrent DVT/PE in adults, following appropriate initial parenteral anticoagulant therapy where required.
- **Prevention of venous thromboembolic events (VTE)** following elective total hip or knee replacement surgery, according to the approved dosing regimen and product labeling. Medicines

Dosage and Administration

- **Recommended Dose:** The appropriate dabigatran dose depends on the indication, age, renal function, bleeding risk and concomitant medicines. **110 mg twice daily** is used in selected adult

patients when dose reduction is appropriate; it should not automatically be considered the standard dose for every indication.
- **Administration:** Swallow the capsule **whole with a glass of water**. It may be taken with or without food.
- **Important: Do not open, chew or crush the capsule**, as this can increase dabigatran exposure and may increase bleeding risk. Medicines
- **Renal Function:** Creatinine clearance (CrCl) should be assessed before treatment and periodically during therapy, particularly in elderly patients or when renal function may deteriorate.
- **Missed Dose:** Follow the physician's instructions; a missed dose should not be doubled. Medicines

Precautions

- **Bleeding Risk:** Dabigatran increases the risk of bleeding. Patients should be monitored for unusual bruising, prolonged bleeding, blood in urine or stool, or unexplained weakness.
- **Renal Impairment:** Dabigatran is predominantly eliminated through the kidneys. Renal function must be assessed before and during treatment. Severe renal impairment is a contraindication in adults. Medicines
- **Elderly Patients:** Bleeding risk may be increased in older patients; dose selection should consider age, renal function and other bleeding-risk factors.
- **Gastrointestinal Disorders:** Use cautiously in patients with gastritis, oesophagitis, gastroesophageal reflux or a history of gastrointestinal bleeding.
- **Surgery & Procedures:** Inform the physician or surgeon before any surgical or invasive procedure because temporary discontinuation may be necessary.
- **Pregnancy & Lactation:** Use during pregnancy or breastfeeding only when specifically considered appropriate by the treating physician.
- **Liver Disease:** Use is not recommended in significant hepatic impairment or liver disease expected to affect survival. Medicines

Contraindications

- Known hypersensitivity to **dabigatran etexilate** or any component of the formulation.
- **Severe renal impairment (CrCl <30 mL/min)** in adults.
- Active clinically significant bleeding.
- Lesions or conditions associated with a significant risk of major bleeding.
- Significant hepatic impairment or liver disease expected to affect survival.
- Concomitant treatment with certain strong **P-glycoprotein (P-gp) inhibitors**, including systemic ketoconazole, cyclosporine, itraconazole and dronedarone.
- Concomitant anticoagulant therapy, except in specific circumstances such as appropriately managed switching of anticoagulants.
- **Prosthetic heart valves requiring anticoagulant treatment.** Medicines

Adverse Reactions

Common:

- Easy bruising

- Nosebleed
- Gastrointestinal discomfort
- Dyspepsia
- Abdominal pain
- Anaemia
- Gastrointestinal bleeding
- Other minor bleeding events

Rare but Serious:

- Major or life-threatening bleeding
- Gastrointestinal haemorrhage
- Intracranial haemorrhage
- Severe allergic/hypersensitivity reactions
- Acute renal deterioration associated with excessive drug exposure or bleeding

Patients should seek immediate medical attention for uncontrolled bleeding, vomiting blood, black/tarry stools, severe headache, weakness or other signs suggestive of serious bleeding. Medicines

Drug Interactions

- **Anticoagulants:** Concurrent use with warfarin, heparins, enoxaparin, rivaroxaban, apixaban or other anticoagulants can substantially increase bleeding risk, except during medically supervised switching.
- **Antiplatelet Agents:** Aspirin and clopidogrel may increase bleeding risk.
- **NSAIDs:** Ibuprofen, diclofenac, ketoprofen and other NSAIDs may increase bleeding risk.
- **P-gp Inhibitors:** Amiodarone, verapamil and quinidine can increase dabigatran exposure; dose selection and monitoring may be required.
- **Strong P-gp Inhibitors:** Ketoconazole, cyclosporine, itraconazole and dronedarone are contraindicated with dabigatran in the referenced product information.
- **P-gp Inducers:** Rifampicin, carbamazepine, phenytoin and St. John's wort may reduce dabigatran concentrations and should generally be avoided.
- Inform the physician about all prescription, OTC and herbal medicines being taken. Medicines

Storage

- Store in a **cool, dry place**, protected from moisture and excessive heat.
- Keep capsules in their original packaging.
- Do not remove capsules from the blister until immediately before use.
- Keep out of reach of children.
- Follow the specific storage conditions stated on the approved product packaging.

Note

This information is intended as a **professional pharmaceutical-literature guideline**. **Dabiflick 110 should be used strictly under the supervision of a qualified healthcare professional.**

Dabigatran dosing is highly dependent on the indication, renal function, age, bleeding risk and concomitant medicines; therefore, the **110 mg strength should not be presented as a universal twice-daily dose** without reference to the approved product indication and prescribing information. Renal function should be assessed before initiation and periodically thereafter. Medicines

Manufactured in India for:

Cafoli Lifecare Pvt. Ltd.
(An ISO 9001: 2015 Certified Co.)
Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113
Corporate Office :
Plot no. 307, Industrial Area Phase-2,
Panchkula, Haryana -134113
TM: Trademark Applied for