Information For the Use of
Dapagliflozin (10mg) + Metformin (1000mg) + Vildagliptin (100mg) Tablet

Dapoflit VM 10/100/1000 Tablet

Dapoflit VM 10/100/1000 Tablet

Composition:

Each tablet contains:

- **Dapagliflozin** – 10 mg
- **Vildagliptin** – 100 mg
- **Metformin Hydrochloride** – 1000 mg

Pharmacological Action:

Dapoflit VM 10/100/1000 Tablet combines three complementary antidiabetic agents that act through different mechanisms to improve glycemic control in patients with **Type 2 Diabetes Mellitus**.

- **SGLT2 Inhibition:** Dapagliflozin selectively inhibits sodium-glucose cotransporter-2 (SGLT2) in the renal proximal tubules, reducing glucose reabsorption and increasing urinary glucose excretion.
- **DPP-4 Inhibition:** Vildagliptin inhibits dipeptidyl peptidase-4 (DPP-4), increasing the activity of incretin hormones and enhancing glucose-dependent insulin secretion while reducing glucagon secretion.
- **Biguanide Action:** Metformin decreases hepatic glucose production, reduces intestinal glucose absorption, and improves peripheral insulin sensitivity.
- **Complementary Glycemic Control:** The combination improves fasting and postprandial glucose control through complementary mechanisms.

Indications:

Dapoflit VM 10/100/1000 Tablet is indicated as an adjunct to **diet and exercise** for the management of:

- **Type 2 Diabetes Mellitus**
- **Inadequately controlled Type 2 Diabetes Mellitus** with combination therapy
- Patients requiring additional glycemic control despite appropriate lifestyle modification and existing antidiabetic treatment

Dosage and Administration:

- **Recommended Dose:** One tablet once daily or as directed by the physician, according to the patient's glycemic response and renal function.
- **Administration:** Take with food to reduce gastrointestinal discomfort associated with metformin.
- Swallow the tablet whole with water; do not crush or chew unless the specific formulation permits it.
- Maintain adequate fluid intake during treatment.
- Continue appropriate diet and exercise as advised by the physician.

Precautions:

- Assess **renal function** before initiation and periodically during treatment.
- Monitor blood glucose and HbA1c regularly.
- Dapagliflozin may cause **volume depletion, hypotension, genital infections, and urinary tract infections**.
- Maintain adequate hydration, particularly in elderly patients or those receiving diuretics.
- Temporarily discontinue treatment and seek medical advice if symptoms suggest **ketoacidosis**, such as nausea, vomiting, abdominal pain, unusual tiredness, or difficulty breathing.
- Metformin carries a rare risk of **lactic acidosis**, particularly in severe renal impairment or conditions associated with tissue hypoxia.
- Vildagliptin should be used cautiously in patients with hepatic impairment; liver function monitoring may be appropriate.
- **Pregnancy and Lactation:** Use only when specifically advised by a physician.

Contraindications:

- Known hypersensitivity to dapagliflozin, metformin, vildagliptin, or any component of the formulation.
- **Severe renal impairment** or dialysis.
- **Metabolic acidosis**, including diabetic ketoacidosis.
- Acute or chronic conditions associated with tissue hypoxia or significant risk of lactic acidosis.
- Severe hepatic impairment where vildagliptin is not appropriate.
- Temporary discontinuation may be required around certain **iodinated contrast imaging procedures**, depending on renal function and clinical circumstances.

Adverse Reactions:

Common:

- Nausea

- Diarrhea
- Abdominal discomfort
- Headache
- Increased urination
- Genital fungal infections
- Urinary tract infections

Less Common but Serious:

- **Lactic acidosis** (rare)
- **Diabetic ketoacidosis**, including cases with relatively modest blood glucose elevation
- Dehydration and hypotension
- Acute kidney injury
- Pancreatitis
- Severe hypersensitivity reactions
- Liver enzyme elevations

Patients should seek immediate medical attention for severe abdominal pain, persistent vomiting, difficulty breathing, severe weakness, or symptoms of dehydration.

Drug Interactions:

- Concomitant use with **insulin or insulin secretagogues** may increase the risk of hypoglycemia.
- Diuretics may increase the risk of **volume depletion and hypotension** with dapagliflozin.
- Alcohol may increase the risk of **metformin-associated lactic acidosis**, particularly with excessive or acute intake.
- Drugs that affect renal function may alter metformin exposure and require monitoring.
- **Iodinated contrast agents** may require temporary interruption of metformin depending on renal function and the imaging procedure.
- Other medicines affecting blood glucose may require dose adjustment by the physician.

Storage:

- Store in a cool, dry place below **25°C**.
- Protect from moisture and direct sunlight.
- Keep out of reach of children.

Note:

This information is intended as a general prescribing guide. Dapoflit VM 10/100/1000 Tablet should be used under the supervision of a registered medical practitioner. The dosage and

duration should be individualized according to the patient's glycemic control, renal function, hepatic status, concomitant medicines, and overall clinical condition.

Because Dapoflit VM combines three glucose-lowering medicines, safe use also depends on checking kidney function, liver enzymes, and sick-day or surgery precautions before and during treatment.

Manufactured in India for:

Cafoli™
LIFECARE

Cafoli Lifecare Pvt. Ltd.
(An ISO 9001: 2015 Certified Co.)
Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113
Corporate Office :
Plot no. 307, Industrial Area Phase-2,
Panchkula, Haryana -134113
TM: Trademark Applied for