Information For the Use of
Dapagliflozin (10 mg) + Teneligliptin (20 mg) + Metformin Hydrochloride Extended Release (500 mg) Tablet

Tengroup DM 20/10/500 Tablet

Tengroup DM 20/10/500 Tablet

Composition:

Each extended-release tablet contains:

- **Teneligliptin** – 20 mg
- **Dapagliflozin** – 10 mg
- **Metformin Hydrochloride Extended Release** – 500 mg

Pharmacological Action:

Tengroup DM 20/10/500 Tablet combines three antidiabetic agents with complementary mechanisms to improve glycemic control in patients with **Type 2 Diabetes Mellitus**.

- **DPP-4 Inhibition:** Teneligliptin inhibits DPP-4, enhancing incretin activity and promoting glucose-dependent insulin secretion while reducing glucagon secretion.
- **SGLT2 Inhibition:** Dapagliflozin inhibits SGLT2 in the renal proximal tubules, reducing glucose reabsorption and increasing urinary glucose excretion.
- **Biguanide Action:** Metformin reduces hepatic glucose production and improves peripheral insulin sensitivity.
- **Complementary Glycemic Control:** The combination acts through different pathways to improve fasting and postprandial blood glucose control.

Indications:

Tengroup DM 20/10/500 Tablet is indicated as an adjunct to **diet and exercise** for the management of:

- **Type 2 Diabetes Mellitus**
- **Inadequately controlled Type 2 Diabetes Mellitus** requiring combination therapy
- Patients requiring additional glycemic control despite appropriate lifestyle modification and existing antidiabetic therapy

---

Dosage and Administration:

- **Recommended Dose:** One tablet once daily or as directed by the physician, according to glycemic response and renal function.
- **Administration:** Take with food to reduce gastrointestinal discomfort associated with metformin.
- Swallow the extended-release tablet whole with water. **Do not crush or chew** unless specifically permitted by the approved formulation.
- Maintain adequate fluid intake during treatment.
- Continue appropriate diet and exercise as advised by the physician.

Precautions:

- Assess **renal function** before initiation and periodically during treatment.
- Monitor blood glucose and HbA1c regularly.
- Dapagliflozin may cause **volume depletion, hypotension, genital infections, and urinary tract infections**.
- Maintain adequate hydration, particularly in elderly patients or those receiving diuretics.
- Metformin carries a rare risk of **lactic acidosis**, especially in severe renal impairment or conditions associated with tissue hypoxia.
- Temporarily discontinue treatment and seek medical advice if symptoms suggest **ketoacidosis**, such as nausea, vomiting, abdominal pain, unusual tiredness, or difficulty breathing.
- Teneligliptin should be used cautiously in patients with hepatic or renal impairment.
- **Pregnancy and Lactation:** Use only when specifically advised by a physician.

Contraindications:

- Known hypersensitivity to dapagliflozin, teneligliptin, metformin, or any component of the formulation.
- **Severe renal impairment** or dialysis.
- **Diabetic ketoacidosis** or metabolic acidosis.
- Acute or chronic conditions associated with significant tissue hypoxia or increased risk of lactic acidosis.
- Severe hepatic impairment where treatment is not considered appropriate.

Adverse Reactions:

Common:

- Nausea
- Diarrhea
- Abdominal discomfort
- Headache

- Increased urination
- Genital fungal infections
- Urinary tract infections
- Dizziness

Less Common but Serious:

- **Lactic acidosis** (rare)
- **Diabetic ketoacidosis**
- Dehydration and hypotension
- Acute kidney injury
- Pancreatitis
- Severe hypersensitivity reactions
- Liver enzyme abnormalities

Patients should seek immediate medical attention for severe abdominal pain, persistent vomiting, difficulty breathing, severe weakness, or symptoms of significant dehydration.

Drug Interactions:

- Concomitant use with **insulin or insulin secretagogues** may increase the risk of hypoglycemia.
- Diuretics may increase the risk of **volume depletion and hypotension** with dapagliflozin.
- Alcohol may increase the risk of **metformin-associated lactic acidosis**, particularly with excessive or acute intake.
- Drugs that affect renal function may alter metformin exposure and require monitoring.
- **Iodinated contrast agents** may require temporary interruption of metformin depending on renal function and the imaging procedure.
- Other medicines affecting blood glucose may require dose adjustment by the physician.

Storage:

- Store in a cool, dry place below **25°C**.
- Protect from moisture and direct sunlight.
- Keep out of reach of children.

Note:

This information is intended as a general prescribing guide. Tengroup DM 20/10/500 Tablet should be used under the supervision of a registered medical practitioner. The dosage and duration should be individualized according to the patient's glycemic control, renal function, hepatic status, concomitant medicines, and overall clinical condition.

Because this fixed-dose combination depends on renal function and the approved metformin release specification, verify the renal-use restrictions and extended-release labeling before finalizing the product literature to avoid inaccurate dosing and contraindication statements.

Manufactured in India for:

Cafoli Lifecare Pvt. Ltd.
(An ISO 9001: 2015 Certified Co.)
Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113
Corporate Office :
Plot no. 307, Industrial Area Phase-2,
Panchkula, Haryana -134113
TM: Trademark Applied for