Dapoflit GM 10/1/500 Tablet

Dapoflit GM 10/1/500 Tablet

Composition:

Each extended-release tablet contains:

- **Dapagliflozin** – 10 mg
- **Glimepiride** – 1 mg
- **Metformin Hydrochloride Extended Release** – 500 mg

Pharmacological Action:

Dapoflit GM 10/1/500 Tablet combines three antidiabetic agents with complementary mechanisms to improve glycemic control in patients with **Type 2 Diabetes Mellitus**.

- **SGLT2 Inhibition:** Dapagliflozin inhibits SGLT2 in the renal proximal tubules, reducing glucose reabsorption and increasing urinary glucose excretion.
- **Insulin Secretagogue Action:** Glimepiride stimulates pancreatic β-cells to release insulin, thereby reducing blood glucose levels.
- **Biguanide Action:** Metformin reduces hepatic glucose production and improves peripheral insulin sensitivity.
- **Complementary Glycemic Control:** The combination targets different pathways involved in glucose regulation, helping improve both fasting and postprandial glycemic control.

Indications:

Dapoflit GM 10/1/500 Tablet is indicated as an adjunct to **diet and exercise** for the management of:

- **Type 2 Diabetes Mellitus**
- **Inadequately controlled Type 2 Diabetes Mellitus** with combination therapy
- Patients requiring additional glycemic control despite appropriate lifestyle modification and existing antidiabetic therapy

---

Dosage and Administration:

- **Recommended Dose:** One tablet once daily or as directed by the physician according to glycemic response and renal function.

- **Administration:** Take with a meal to reduce gastrointestinal discomfort and minimize the risk of hypoglycemia associated with glimepiride.
- Swallow the extended-release tablet whole with water. **Do not crush, chew, or split** unless specifically permitted for the particular formulation.
- Maintain adequate fluid intake during treatment.
- Continue appropriate diet and exercise as advised by the physician.

Precautions:

- Assess **renal function** before initiation and periodically during treatment.
- Monitor blood glucose and HbA1c regularly.
- Glimepiride may cause **hypoglycemia**; patients should be educated about recognizing symptoms such as sweating, tremor, dizziness, hunger, and confusion.
- Dapagliflozin may cause **volume depletion, hypotension, genital infections, and urinary tract infections**.
- Maintain adequate hydration, particularly in elderly patients or those taking diuretics.
- Metformin carries a rare risk of **lactic acidosis**, especially in severe renal impairment or conditions associated with tissue hypoxia.
- Temporarily interrupt treatment and seek medical advice if symptoms suggest **ketoacidosis**, such as nausea, vomiting, abdominal pain, unusual fatigue, or difficulty breathing.
- **Pregnancy and Lactation:** Use only when specifically advised by a physician.

Contraindications:

- Known hypersensitivity to dapagliflozin, glimepiride, metformin, or any component of the formulation.
- **Severe renal impairment** or dialysis.
- **Diabetic ketoacidosis** or metabolic acidosis.
- Acute or chronic conditions associated with significant tissue hypoxia or increased risk of lactic acidosis.
- Severe hepatic impairment where glimepiride or metformin is not appropriate.
- Temporary discontinuation may be required around certain **iodinated contrast imaging procedures**, depending on renal function and clinical circumstances.

Adverse Reactions:

Common:

- Hypoglycemia
- Nausea
- Diarrhea
- Abdominal discomfort
- Headache

- Increased urination
- Genital fungal infections
- Urinary tract infections

Less Common but Serious:

- **Lactic acidosis** (rare)
- **Diabetic ketoacidosis**
- Severe hypoglycemia
- Dehydration and hypotension
- Acute kidney injury
- Severe hypersensitivity reactions

Patients should seek immediate medical attention for severe or persistent hypoglycemia, difficulty breathing, severe abdominal pain, persistent vomiting, confusion, or marked weakness.

Drug Interactions:

- Other **antidiabetic medicines**, particularly insulin or insulin secretagogues, may increase the risk of hypoglycemia.
- Diuretics may increase the risk of **volume depletion and hypotension** with dapagliflozin.
- Alcohol may increase the risk of hypoglycemia and may also increase the risk of **metformin-associated lactic acidosis**, particularly with excessive or acute intake.
- Drugs that impair renal function may increase metformin exposure and require monitoring.
- **Iodinated contrast agents** may require temporary interruption of metformin depending on renal function and the imaging procedure.
- Beta-blockers may mask some symptoms of hypoglycemia.

Storage:

- Store in a cool, dry place below **25°C**.
- Protect from moisture and direct sunlight.
- Keep out of reach of children.

Note:

This information is intended as a general prescribing guide. Dapoflit GM 10/1/500 Tablet should be used under the supervision of a registered medical practitioner. The dosage and duration should be individualized according to the patient's glycemic control, renal function, hepatic status, risk of hypoglycemia, concomitant medicines, and overall clinical condition.

Because glimepiride adds a meaningful risk of hypoglycemia to Dapoflit GM, safe use depends on recognizing low-blood-sugar symptoms and knowing how to respond while taking the tablet.

Manufactured in India for:

Cafoli Lifecare Pvt. Ltd.
(An ISO 9001: 2015 Certified Co.)
Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113
Corporate Office :
Plot no. 307, Industrial Area Phase-2,
Panchkula, Haryana -134113
TM: Trademark Applied for