Information For the Use of
Drotaverine (40mg) Tablets

Dotahub 40 Tablet

DOTAHUB 40 TABLET

Drotaverine (40 mg)

1. Composition

Each uncoated tablet contains:
Drotaverine Hydrochloride equivalent to Drotaverine 40 mg
Excipients q.s.

2. Pharmacological Action

Drotaverine is a **smooth-muscle antispasmodic**. It primarily inhibits phosphodiesterase-4 (PDE-4), resulting in increased intracellular cyclic AMP and relaxation of smooth muscle.

It helps relieve pain and discomfort associated with **smooth-muscle spasms** of the gastrointestinal, biliary and urinary tracts.

3. Indications

Drotaverine is indicated for the symptomatic relief of pain associated with smooth-muscle spasm, including:

- **Abdominal cramps and intestinal spasms**
- Biliary tract spasms and colicky pain
- Renal/urinary tract spasms
- Dysmenorrhoea and uterine spasmodic pain
- Other conditions associated with visceral smooth-muscle spasm

4. Dosage and Administration

Adults: The usual oral dose is **40–80 mg, 2–3 times daily**, or as directed by the physician.

Swallow the tablet with water. It may be taken with or without food.

The dose and duration should be individualized according to the severity and cause of symptoms.

5. Precautions

- Use cautiously in patients with **severe hepatic, renal or cardiac impairment**.
- Patients with low blood pressure should use drotaverine with caution, as hypotension may occur.
- Dizziness may occur; caution is advised while driving or operating machinery if affected.

- Persistent or unexplained abdominal pain should be medically evaluated rather than repeatedly treated with an antispasmodic.
- Use during pregnancy and lactation should be based on medical advice.

6. Contraindications

Drotaverine is contraindicated in patients with:

- Known hypersensitivity to **drotaverine or any component of the formulation**.
- Severe hepatic impairment.
- Severe renal impairment.
- Severe cardiac insufficiency.

7. Adverse Reactions

Possible adverse reactions include:

- Nausea
- Constipation
- Headache
- Dizziness or vertigo
- Palpitations
- Decreased blood pressure

Rarely, hypersensitivity reactions such as rash, itching or angioedema may occur.

8. Drug Interactions

- Concomitant use with other medicines that lower blood pressure may increase the risk of **hypotension**.
- Drotaverine may interact with **levodopa**, potentially reducing its antiparkinsonian effect and worsening rigidity or tremor.
- Inform the physician about all prescription, OTC and herbal medicines being used.

9. Storage

Store below **30°C**, protected from light and moisture.

Keep out of reach of children.

10. Note / Important

Drotaverine provides **symptomatic relief of smooth-muscle spasms** and does not treat the underlying cause of persistent abdominal or pelvic pain.

If pain is severe, recurrent, unexplained, or associated with fever, vomiting, bleeding or other concerning symptoms, appropriate medical evaluation is required.

For use under medical supervision.

Keep out of reach of children.

Manufactured in India for:

Cafoli Lifecare Pvt. Ltd.
(An ISO 9001: 2015 Certified Co.)
Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113
Corporate Office :
Plot no. 307, Industrial Area Phase-2,
Panchkula, Haryana -134113
TM: Trademark Applied for