Information For the Use of
Doxylamine (20mg) + Vitamin B6 (Pyridoxine) (20mg) + Folic Acid (5mg) Tablet

Doxaloom Forte Tablet

Brand Name: Doxaloom Forte Tablet
Dosage Form: Tablet
Composition:
Each film-coated tablet contains:
• Doxylamine Succinate IP 20 mg
• Pyridoxine Hydrochloride IP (Vitamin B6) 20 mg
• Folic Acid IP 5 mg

Description:
Doxaloom Forte Tablet is a fixed-dose combination of Doxylamine Succinate, an antihistamine with antiemetic properties; Pyridoxine (Vitamin B6), an essential vitamin involved in metabolic functions; and Folic Acid, a vitamin required for normal fetal development and red blood cell formation. It is indicated for the management of nausea and vomiting associated with pregnancy and for providing folate supplementation during pregnancy. It helps in reducing pregnancy-related nausea while supporting maternal nutritional requirements.

Indications:
Indicated for the treatment and management of:
• Nausea and vomiting during pregnancy (morning sickness)
• Pregnancy-associated nutritional supplementation
• Folic acid deficiency during pregnancy
• Vitamin B6 supplementation as advised by the physician
It helps in relieving nausea and vomiting while supporting healthy maternal and fetal nutritional status.

Dosage & Administration:
Dosage should be as directed by the physician.
The tablet should be swallowed whole with water, preferably at bedtime or as prescribed.
Do not exceed the recommended dosage.

Pharmacological Action:

- **Doxylamine Succinate is an H1-histamine receptor antagonist that exerts antiemetic effects by suppressing the vomiting center, thereby reducing nausea and vomiting.**
- **Pyridoxine (Vitamin B6) plays a vital role in amino acid metabolism, neurotransmitter synthesis, and contributes to the management of pregnancy-related nausea.**
- **Folic Acid is essential for DNA synthesis, cell division, and normal fetal neural tube development while supporting erythropoiesis.**

Together, these mechanisms provide effective relief from pregnancy-related nausea and support maternal nutritional health.

Side Effects:
Common side effects include drowsiness, dizziness, dry mouth, constipation, fatigue, and blurred vision.
Occasionally, headache or gastrointestinal discomfort may occur.
Rarely, hypersensitivity reactions may be observed.

Precautions & Warnings:

- **Use only under the supervision of a qualified healthcare professional during pregnancy**
- **May cause drowsiness; avoid driving or operating machinery if affected**
- **Avoid alcohol and other central nervous system depressants during treatment**
- **Use with caution in patients with glaucoma, asthma, urinary retention, or hepatic impairment**
- **Do not exceed the prescribed dosage**

Drug Interactions:

Concurrent use with alcohol, sedatives, tranquilizers, opioids, or other central nervous system depressants may enhance drowsiness.
Use cautiously with anticholinergic medications due to the potential for additive effects.

Storage:

Store below 25°C in a cool, dry place.
Protect from light and moisture.
Keep out of reach of children.

Manufactured in India for:

Cafoli™
LIFECARE

Cafoli Lifecare Pvt. Ltd.
(An ISO 9001: 2015 Certified Co.)
Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113
Corporate Office :
Plot no. 307, Industrial Area Phase-2,
Panchkula, Haryana -134113
TM: Trademark Applied for