Information For the Use of
Esomeprazole (40mg) + Levosulpiride (75mg) Capsule

Esoable L Capsule

Brand Name: Esoable L Capsule
Dosage Form: Capsule
Composition:
Each capsule contains:
• Esomeprazole Magnesium IP equivalent to Esomeprazole 40 mg
• Levosulpiride SR IP 75 mg

Description:
Esoable L Capsule is a fixed-dose combination of Esomeprazole, a proton pump inhibitor (PPI), and Levosulpiride, a prokinetic agent. It is indicated for the management of gastroesophageal reflux disease (GERD), dyspepsia, and acid-related gastrointestinal disorders associated with impaired gastric motility. It helps in reducing gastric acid secretion, improving gastric emptying, and relieving symptoms such as heartburn, acid regurgitation, bloating, and upper abdominal discomfort.

Indications:
Indicated for the treatment of:
• Gastroesophageal reflux disease (GERD)
• Acid peptic disorders
• Functional dyspepsia
• Gastritis associated with acid secretion
• Reflux symptoms with delayed gastric emptying
It helps in relieving heartburn, acid reflux, bloating, nausea, and indigestion while promoting gastrointestinal comfort.

Dosage & Administration:
Dosage should be as directed by the physician.
The capsule should preferably be taken before meals, usually in the morning.
Swallow the capsule whole without crushing or chewing.
Follow the prescribed treatment duration for optimal therapeutic benefit.

Pharmacological Action:

- **Esomeprazole irreversibly inhibits the H+/K+-ATPase enzyme system (proton pump) in gastric parietal cells, thereby suppressing gastric acid secretion.**
- **Levosulpiride enhances gastrointestinal motility by selectively blocking dopamine D2 receptors, promoting gastric emptying and reducing reflux symptoms.**

Together, these mechanisms provide effective acid suppression and improvement in gastrointestinal motility.

Side Effects:
Common side effects include headache, nausea, abdominal pain, constipation, diarrhea, dizziness, and dry mouth.
Occasionally, fatigue, sleep disturbances, or gastrointestinal discomfort may occur.
Rarely, prolonged use may be associated with vitamin B12 deficiency or electrolyte imbalance.

Precautions & Warnings:

- **Use with caution in patients with hepatic impairment**
- **Long-term use should be monitored by a healthcare professional**
- **Rule out gastric malignancy before initiating prolonged therapy**
- **Use cautiously in elderly patients**
- **Pregnancy and lactation use should be based on physician advice**

Drug Interactions:
May interact with anticoagulants, antiplatelet agents, digoxin, ketoconazole, and other medications affected by gastric pH.
Concurrent use with central nervous system-active drugs may require monitoring due to Levosulpiride.

Storage:
Store below 25°C in a cool, dry place.
Protect from light and moisture.
Keep out of reach of children.

Manufactured in India for:

Cafoli™
LIFECARE

Cafoli Lifecare Pvt. Ltd.
(An ISO 9001: 2015 Certified Co.)
Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113
Corporate Office :
Plot no. 307, Industrial Area Phase-2,
Panchkula, Haryana -134113
TM: Trademark Applied for