Information For the Use of
Fenofibrate (200mg) Capsule

Fenoflick 200 Capsule

Brand Name: Fenoflick 200 Capsule
Dosage Form: Capsule
Composition:
Each capsule contains:
• Fenofibrate IP 200 mg

Description:
Fenoflick 200 Capsule contains Fenofibrate, a fibric acid derivative indicated for the management of dyslipidemia and hypertriglyceridemia. Fenofibrate acts as a peroxisome proliferator-activated receptor alpha (PPAR-α) agonist, improving lipid metabolism by reducing serum triglycerides, lowering low-density lipoprotein (LDL) cholesterol, and increasing high-density lipoprotein (HDL) cholesterol. It is used as an adjunct to diet, exercise, and lifestyle modification to reduce cardiovascular risk in appropriate patients.

Indications:
Indicated for the treatment of:
• Primary hypercholesterolemia
• Mixed dyslipidemia
• Hypertriglyceridemia
• Diabetic dyslipidemia
• Prevention of pancreatitis associated with severe hypertriglyceridemia
• Adjunct to diet and lifestyle modification for lipid control
It helps reduce triglyceride and LDL cholesterol levels while increasing HDL cholesterol, thereby improving the overall lipid profile.

Dosage & Administration:
Dosage should be as directed by the physician.
The capsule should be swallowed whole with water, preferably with meals to enhance absorption.
Patients should continue a lipid-lowering diet and lifestyle modifications during therapy.
Periodic monitoring of lipid profile, liver function, and renal function is recommended.

Pharmacological Action:

- **Fenofibrate is a PPAR-α agonist that enhances lipolysis and the elimination of triglyceride-rich lipoproteins by activating lipoprotein lipase.**
- **It reduces hepatic production of very low-density lipoprotein (VLDL), lowers triglycerides and LDL cholesterol, and increases HDL cholesterol by promoting apolipoprotein A-I and A-II synthesis.**
- **These actions contribute to improved lipid metabolism and reduced cardiovascular risk in patients with dyslipidemia.**

Side Effects:
Common side effects include nausea, abdominal pain, dyspepsia, diarrhea, headache, back pain, and elevated liver enzymes.
Rarely, myopathy, rhabdomyolysis, gallstones, pancreatitis, hypersensitivity reactions, or hepatic dysfunction may occur.

Precautions & Warnings:

• Monitor lipid profile, liver function tests, and renal function periodically during treatment.

• Use with caution in patients with hepatic impairment, renal impairment, gallbladder disease, or hypothyroidism.

• Patients should report unexplained muscle pain, tenderness, or weakness immediately, especially when used with statins.

• Contraindicated in patients with severe hepatic disease, severe renal impairment, active gallbladder disease, or known hypersensitivity to Fenofibrate.

• Use during pregnancy and lactation only if the potential benefit justifies the potential risk.

Drug Interactions:

May interact with statins, oral anticoagulants (e.g., warfarin), cyclosporine, bile acid sequestrants, and other lipid-lowering agents. Concomitant use may increase the risk of myopathy or alter anticoagulant effects; appropriate monitoring and dose adjustment may be required.

Storage:

Store below 25°C in a cool, dry place.

Protect from light and moisture.

Keep out of reach of children.

Manufactured in India for:

Cafoli™
LIFECARE

Cafoli Lifecare Pvt. Ltd.

(An ISO 9001: 2015 Certified Co.)

Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113

Corporate Office :

Plot no. 307, Industrial Area Phase-2,
Panchkula, Haryana -134113

TM: Trademark Applied for