Information For the Use of
Finasteride (1mg) Tablets

Finaslide 1 Tablet

FINASLIDE 1 TABLET

Finasteride (1 mg)

1. Composition

Each uncoated tablet contains:
Finasteride IP 1 mg
Excipients q.s.

2. Pharmacological Action

Finasteride is a specific inhibitor of **Type II 5-alpha-reductase**, the enzyme responsible for converting testosterone into dihydrotestosterone (DHT). By reducing DHT levels in the scalp, finasteride helps reduce follicular miniaturization and may promote the maintenance and regrowth of hair in men with androgenetic alopecia.

3. Indications

Finasteride 1 mg is indicated for the treatment of **male pattern hair loss (androgenetic alopecia)** in adult men.

It may help:

- Reduce further hair loss.
- Promote hair regrowth, particularly in the vertex and frontal/mid-scalp regions.
- Maintain existing hair growth with continued treatment.

4. Dosage and Administration

Recommended dose: One tablet of **Finasteride 1 mg once daily**, with or without food.

The tablet should be swallowed whole with water and should not be crushed or broken.

Continuous treatment is generally required to maintain the therapeutic benefit. Improvement may take several months of regular treatment.

5. Precautions

- Finasteride is intended for **use in men** for this indication.
- Women who are pregnant or may become pregnant should **not handle crushed or broken tablets**, because finasteride may cause abnormalities of the external genitalia in a male fetus.

- Finasteride may reduce serum PSA levels; inform the physician before PSA testing.
- Assess patients with significant urinary symptoms for other underlying causes before treatment.
- Caution is advised in patients with hepatic impairment, as finasteride is extensively metabolized in the liver.
- Report persistent sexual, mood-related, breast-related, or other unusual symptoms to the physician.

6. Contraindications

Finasteride is contraindicated in:

- Patients with known hypersensitivity to finasteride or any component of the formulation.
- **Women who are pregnant or may become pregnant.**
- **Children and adolescents.**

7. Adverse Reactions

Possible adverse effects include:

- Decreased libido
- Erectile dysfunction
- Ejaculatory disorders / decreased volume of ejaculate
- Breast tenderness or enlargement
- Rash or hypersensitivity reactions

Post-marketing reports have also included persistent sexual dysfunction and mood-related symptoms in some patients. Patients should seek medical advice if such symptoms occur.

8. Drug Interactions

Clinically significant drug interactions are uncommon with finasteride. However:

- Inform the physician about all prescription, OTC and herbal medicines being used.
- Finasteride may alter PSA values, which should be considered when interpreting PSA results.
- No clinically important interaction has been established with commonly used antihypertensive, antidiabetic or NSAID medicines.

9. Storage

Store below **30°C**, protected from light and moisture.

Keep out of reach of children.

10. Note / Important

Finasteride 1 mg is specifically used for **male androgenetic alopecia**. It does not provide an immediate effect; noticeable improvement generally requires several months of continuous treatment.

Treatment should be continued as advised by the physician to maintain the benefit. Women who are pregnant or may become pregnant should not handle crushed or broken finasteride tablets.

For use under medical supervision.

Keep out of reach of children.

Manufactured in India for:

Cafoli™
LIFECARE

Cafoli Lifecare Pvt. Ltd.
(An ISO 9001: 2015 Certified Co.)
Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113
Corporate Office :
Plot no. 307, Industrial Area Phase-2,
Panchkula, Haryana -134113
TM: Trademark Applied for