Information For the Use of
Cefpodoxime Proxetil (100mg) Tablet

Flickdox 100 DT Tablet

Brand Name: Flickdox 100 DT Tablet
Dosage Form: Dispersible Tablet (DT)
Composition:
Each dispersible tablet contains:
• Cefpodoxime Proxetil IP equivalent to Cefpodoxime 100 mg

Description:
Flickdox 100 DT Tablet contains Cefpodoxime Proxetil, an orally administered third-generation cephalosporin antibiotic. It is indicated for the treatment of mild to moderate bacterial infections caused by susceptible microorganisms. Cefpodoxime Proxetil is a prodrug that is converted to its active form, Cefpodoxime, after absorption. It exerts broad-spectrum bactericidal activity against both Gram-positive and Gram-negative bacteria.

Indications:
Indicated for the treatment of:
• Upper and lower respiratory tract infections
• Acute otitis media
• Pharyngitis and tonsillitis
• Acute bacterial sinusitis
• Urinary tract infections
• Skin and soft tissue infections
• Uncomplicated bacterial infections caused by susceptible organisms
It helps in eliminating susceptible bacteria, relieving infection-related symptoms, and preventing the progression of bacterial infections.

Dosage & Administration:
Dosage should be as directed by the physician.
Disperse the tablet in a small quantity of water before administration or swallow whole if advised.
It is preferably administered with food to enhance absorption.
Complete the full prescribed course of therapy even if symptoms improve earlier.

Pharmacological Action:

- **Cefpodoxime Proxetil is a prodrug that is hydrolyzed in vivo to Cefpodoxime, its active antibacterial form.**
- **Cefpodoxime inhibits bacterial cell wall synthesis by binding to penicillin-binding proteins (PBPs), resulting in bacterial cell lysis and death.**
- **It exhibits broad-spectrum bactericidal activity against susceptible Gram-positive and Gram-negative pathogens.**

Side Effects:
Common side effects include diarrhea, nausea, vomiting, abdominal pain, headache, dizziness, skin rash, and mild gastrointestinal discomfort.
Rarely, hypersensitivity reactions, pseudomembranous colitis, or severe allergic reactions may occur.

Precautions & Warnings:

• Use only for bacterial infections; it is not effective against viral infections
• Use with caution in patients with renal impairment or a history of gastrointestinal disease, especially colitis
• Patients with known hypersensitivity to cephalosporins or penicillins should use with caution
• Prolonged use may result in overgrowth of non-susceptible organisms
• Complete the prescribed course to reduce the risk of antimicrobial resistance.

Drug Interactions:
Concomitant use with antacids or H_2-receptor antagonists may reduce the absorption of Cefpodoxime.
Probenecid may increase serum concentrations of Cefpodoxime.
Use cautiously with nephrotoxic drugs due to the potential for additive renal effects.

Storage:
Store below 25°C in a cool, dry place.
Protect from light and moisture.
Keep out of reach of children.

Manufactured in India for:

Cafoli™
LIFECARE

Cafoli Lifecare Pvt. Ltd.
(An ISO 9001: 2015 Certified Co.)
Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113
Corporate Office :
Plot no. 307, Industrial Area Phase-2,
Panchkula, Haryana -134113
TM: Trademark Applied for