Information For the Use of
Fluorometholone IP 0.1% w/v + Benzalkonium Chloride solution IP 0.02% v/v Eye Drops

Flonshell Eye Drops

Brand Name: Flonshell Eye Drops
Dosage Form: Eye Drops
Composition:
Each ml contains:

- **Fluorometholone IP 0.1% w/v**
- **Benzalkonium Chloride Solution IP 0.02% v/v**
- **Sterile aqueous vehicle q.s.**

Description:
Flonshell Eye Drops contains Fluorometholone, a topical ophthalmic corticosteroid with anti-inflammatory activity. It is used for the treatment of corticosteroid-responsive inflammatory conditions affecting the conjunctiva, cornea, and anterior segment of the eye. Fluorometholone suppresses ocular inflammation and helps reduce redness, swelling, irritation, and associated discomfort.
Indications:
Indicated for the treatment of:

- **Corticosteroid-responsive ocular inflammation.**
- **Inflammatory conditions of the palpebral and bulbar conjunctiva.**
- **Corneal inflammation.**
- **Inflammation of the anterior segment of the eye.**
- **Ocular redness, swelling, and irritation associated with steroid-responsive inflammation.**

Dosage & Administration:
Instill the prescribed number of drops into the affected eye(s) as directed by the ophthalmologist.
Shake well before use if the preparation is a suspension.
Avoid touching the dropper tip to the eye or surrounding surfaces.
If more than one ophthalmic medication is being used, maintain an appropriate interval between applications.
Use only for the duration prescribed by the ophthalmologist.
Pharmacological Action:

- **Fluorometholone is an ophthalmic corticosteroid that suppresses inflammatory responses in ocular tissues.**
- **It reduces inflammatory mediator activity and inhibits processes such as capillary dilation, leukocyte migration, edema, and other components of the inflammatory response.**
- **These actions help reduce ocular redness, swelling, irritation, and inflammation.**

Side Effects:
Common adverse effects may include temporary burning or stinging following instillation and mild ocular irritation.
Prolonged or inappropriate use may cause increased intraocular pressure, glaucoma, delayed wound healing, cataract formation, or secondary ocular infections.
Precautions & Warnings:

- **For ophthalmic use only.**
- **Use under the supervision of an ophthalmologist.**
- **Prolonged use may increase intraocular pressure; monitoring is recommended during extended therapy.**
- **Corticosteroids may mask or worsen certain ocular infections.**
- **Avoid use in untreated fungal, mycobacterial, or certain viral infections of the eye, including**

epithelial herpes simplex keratitis.

- **Use cautiously in patients with glaucoma or a history of steroid-induced elevation of intraocular pressure.**
- **Prolonged corticosteroid therapy may delay corneal healing and increase the risk of secondary infection.**

Drug Interactions:

Concomitant use with other ophthalmic corticosteroids may increase the risk of steroid-related adverse effects.

Patients using multiple ophthalmic preparations should inform their ophthalmologist.

The preservative Benzalkonium Chloride may be absorbed by soft contact lenses; contact lenses should generally be removed before instillation and reinserted as advised.

Storage:

Store below 25°C in a cool, dry place.

Protect from light.

Do not freeze.

Keep the container tightly closed when not in use.

Keep out of reach of children.

Manufactured in India for:

Cafoli™
LIFECARE

Cafoli Lifecare Pvt. Ltd.

(An ISO 9001: 2015 Certified Co.)

Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113

Corporate Office :

Plot no. 307, Industrial Area Phase-2,
Panchkula, Haryana -134113

TM: Trademark Applied for