Information For the Use of
Flubiprofen Sodium IP 0.03% w/v + Hydroxypropylmethylcellulose IP 0.25% w/v + Phenylmercuric Nitrate IP 0.001% w/v Eye Drops

Flublink Plus Eye Drop

Flublink Plus Eye Drops

Composition:

Each mL contains:

- **Flurbiprofen Sodium IP** – 0.03% w/v
- **Hydroxypropylmethylcellulose IP** – 0.25% w/v
- **Phenylmercuric Nitrate IP** – 0.001% w/v

Pharmacological Action:

Flublink Plus Eye Drops combines a **topical non-steroidal anti-inflammatory drug (NSAID)** with a **lubricating/viscoelastic agent**.

- **Anti-inflammatory Action:** Flurbiprofen inhibits cyclooxygenase enzymes, reducing prostaglandin synthesis and thereby reducing ocular inflammation and pain.
- **Analgesic Action:** Reduction of prostaglandin-mediated inflammation helps relieve ocular discomfort.
- **Lubricating Action:** Hydroxypropylmethylcellulose forms a protective film over the ocular surface and helps maintain moisture.
- **Ocular Surface Protection:** The lubricating component helps reduce irritation and improves ocular comfort.

Indications:

Flublink Plus Eye Drops is indicated for the management of:

- **Post-operative Ocular Inflammation**
- **Ocular Pain and Inflammation following Ophthalmic Procedures**
- **Inflammatory Eye Conditions responsive to topical NSAID therapy**
- **Ocular Surface Irritation associated with inflammation**, when prescribed by an ophthalmologist

---

Dosage and Administration:

- **Recommended Dose:** Instill **1 drop into the affected eye(s) as directed by the ophthalmologist**.

- Wash hands before administration.
- Tilt the head back and gently pull down the lower eyelid.
- Instill the prescribed drop without touching the dropper tip to the eye, eyelids, or surrounding surfaces.
- Close the eye gently after administration.
- If more than one ophthalmic preparation is being used, maintain an appropriate interval between applications.
- Use for the duration prescribed by the ophthalmologist.

Precautions:

- For **ophthalmic use only**.
- Use cautiously in patients with a history of **NSAID or aspirin hypersensitivity**.
- Topical ophthalmic NSAIDs may delay corneal healing in susceptible patients.
- Use cautiously in patients with **corneal epithelial defects, dry eye, corneal disease, or previous ocular surgery**.
- Prolonged or frequent use may increase the risk of **corneal complications**.
- Avoid contamination of the dropper tip.
- Contact lenses should generally be removed before administration and reinserted only as advised by the ophthalmologist.
- **Pregnancy and Lactation:** Use only when clearly indicated and under medical supervision.

Contraindications:

- Known hypersensitivity to **flurbiprofen, other NSAIDs, or any component of the formulation**.
- Previous aspirin/NSAID-induced asthma, urticaria, or other significant hypersensitivity reactions.
- Active or suspected ocular infection where NSAID therapy is inappropriate.
- Use cautiously or avoid in patients with significant corneal epithelial defects unless specifically directed by an ophthalmologist.

Adverse Reactions:

Common:

- Temporary burning or stinging
- Eye irritation
- Temporary blurred vision
- Redness
- Watery eyes

Less Common but Serious:

- Delayed corneal healing
- Corneal epithelial breakdown
- Corneal ulceration or thinning
- Increased ocular irritation
- Hypersensitivity reactions

Seek immediate ophthalmic attention if there is severe eye pain, worsening redness, reduced vision, or persistent irritation.

Drug Interactions:

- Concurrent use with other **topical ophthalmic NSAIDs** may increase the risk of ocular adverse effects.
- Concurrent topical **corticosteroids** may increase the risk of delayed corneal healing and other corneal complications and should be used only under ophthalmic supervision.
- If multiple ophthalmic preparations are prescribed, administer them separately with an appropriate interval.
- Inform the ophthalmologist about systemic NSAIDs, aspirin, anticoagulants, and other medicines being used.

Storage:

- Store in a cool, dry place below **25°C**.
- Protect from direct sunlight and excessive heat.
- Keep the bottle tightly closed when not in use.
- Do not touch the dropper tip to any surface.
- Keep out of reach of children.

Note:

This information is intended as a general prescribing guide. **Flublink Plus Eye Drops should be used under the supervision of a qualified ophthalmologist.** Topical ophthalmic NSAIDs should be used carefully in patients with corneal epithelial defects or ocular surface disease because prolonged or inappropriate use may result in serious corneal complications.

Important: The listed formulation contains **phenylmercuric nitrate 0.001% w/v** as a preservative. The final IP literature should match the approved product specification and current regulatory requirements for the formulation and preservative.

Manufactured in India for:

Cafoli Lifecare Pvt. Ltd.
(An ISO 9001: 2015 Certified Co.)
Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113
Corporate Office :
Plot no. 307, Industrial Area Phase-2,
Panchkula, Haryana -134113
TM: Trademark Applied for