Information For the Use of
Fluticasone Propionate (0.05% w/w) Cream

Flutialfa Cream

FLUTIALFA CREAM

Composition

Each gram of cream contains:

Fluticasone Propionate I.P. 0.05% w/w
Cream base … q.s.

Pharmacological Action

Fluticasone Propionate is a potent topical corticosteroid with anti-inflammatory, antipruritic and vasoconstrictor properties.

It acts by reducing the release and activity of inflammatory mediators and suppressing inflammatory responses in the skin. This results in reduction of **redness, swelling, itching and irritation** associated with corticosteroid-responsive dermatoses.

Indications

Flutialfa Cream is indicated for the relief of inflammatory and pruritic manifestations of corticosteroid-responsive skin conditions, including:

- Atopic dermatitis
- Eczema
- Contact dermatitis
- Seborrhoeic dermatitis
- Psoriasis
- Other steroid-responsive inflammatory dermatoses

It should be used only for conditions responsive to topical corticosteroid therapy.

Dosage and Administration

Adults and children over 1 year:
Apply a **thin layer** of Flutialfa Cream to the affected areas **once or twice daily**, or as directed by the physician.

Gently massage into the affected area until absorbed.

Treatment should generally be limited to the **shortest duration necessary** to control symptoms.

Avoid prolonged continuous treatment, particularly on the face, skin folds or other sensitive areas.

Precautions

- For **external use only**.
- Avoid contact with the eyes and mucous membranes.
- Do not apply to the face, groin or axillae for prolonged periods unless specifically prescribed.
- Avoid application to large areas of the body or under occlusive dressings unless directed by a physician.
- Prolonged or excessive use may cause **skin atrophy, striae and telangiectasia**.
- Use cautiously in children because they may be more susceptible to systemic corticosteroid effects.
- Do not use on untreated bacterial, fungal or viral skin infections.
- If irritation, sensitization or worsening of the condition occurs, treatment should be discontinued and medical advice sought.
- Use cautiously on broken or damaged skin because systemic absorption may be increased.

Contraindications

Flutialfa Cream is contraindicated in:

- Known hypersensitivity to **Fluticasone Propionate** or any component of the formulation.
- Untreated bacterial, fungal or viral infections of the skin.
- Rosacea.
- Acne vulgaris.
- Perioral dermatitis.
- Pruritus without inflammation where topical corticosteroid treatment is inappropriate.

It should not be used for conditions that are not responsive to topical corticosteroid therapy.

Adverse Reactions

Common/Local:

- Burning or stinging
- Itching
- Irritation
- Dryness
- Redness

With prolonged or inappropriate use:

- Skin thinning (atrophy)
- Striae
- Telangiectasia
- Folliculitis
- Acneiform eruptions
- Hypopigmentation
- Contact dermatitis
- Changes in skin pigmentation

Rare/Systemic:

Excessive application, prolonged use, application to large areas or use under occlusion may increase systemic absorption and, rarely, cause **hypothalamic-pituitary-adrenal (HPA) axis suppression** or other systemic corticosteroid effects.

Drug Interactions

Clinically significant interactions are uncommon with appropriate topical use.

However, concomitant use of other **topical corticosteroids or irritating topical preparations** on the same area may increase the risk of local adverse effects.

Patients should inform their physician about all topical and systemic medicines being used.

Pregnancy and Lactation

Use during pregnancy should be considered only when the expected benefit justifies the potential risk.

Avoid prolonged or extensive application during pregnancy unless specifically prescribed.

During lactation, use only on medical advice. **Do not apply to the breast or nipple immediately before breastfeeding.** If treatment is required on the breast, the area should be thoroughly cleansed before feeding.

Storage

Store in a **cool, dry place**, protected from excessive heat and direct sunlight.

Keep the tube tightly closed and **out of reach of children**.

Do not use after the expiry date.

Note

For external use only. For use under medical supervision.

Apply a thin layer only to the affected areas and use for the shortest duration necessary. Avoid prolonged or extensive use unless specifically prescribed.

Important: Product-specific approved prescribing information should take precedence for the final printed literature, particularly regarding age restrictions, treatment duration and application frequency.

Manufactured in India for:

Cafoli Lifecare Pvt. Ltd.
(An ISO 9001: 2015 Certified Co.)
Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113
Corporate Office :
Plot no. 307, Industrial Area Phase-2,
Panchkula, Haryana -134113
TM: Trademark Applied for