Information For the Use of
Formoterol (6mcg) + Fluticasone Propionate (250mcg)_CFC Free Inhaler

Flutialfa F 250 Inhaler

Flutialfa F 250 Inhaler

Composition:

Each metered dose contains:

- **Formoterol** – 6 mcg
- **Fluticasone Propionate** – 250 mcg
- **CFC-Free Propellant**

Pharmacological Action:

Flutialfa F 250 combines a **long-acting β_2-adrenergic agonist (LABA)** with an **inhaled corticosteroid (ICS)** to provide sustained bronchodilation and control of airway inflammation.

- **Bronchodilation:** Formoterol stimulates β_2 receptors in bronchial smooth muscle, producing rapid and prolonged relaxation of the airways.
- **Anti-inflammatory Action:** Fluticasone propionate reduces airway inflammation, edema, and inflammatory mediator activity.
- **Improved Airflow:** The combination reduces bronchoconstriction and helps improve lung function.
- **Maintenance Therapy:** Provides sustained control of airway symptoms when regular inhaled corticosteroid/LABA therapy is indicated.

Indications:

Flutialfa F 250 Inhaler is indicated for the **maintenance treatment** of:

- **Bronchial Asthma**
- **Chronic Obstructive Pulmonary Disease (COPD)**, in appropriately selected patients
- Other chronic obstructive airway conditions where an inhaled corticosteroid/LABA combination is prescribed

Note: This inhaler is intended for **regular maintenance therapy** and should not be relied upon as the sole treatment for sudden, severe bronchospasm unless specifically prescribed as part of an appropriate regimen.

Dosage and Administration:

- **Recommended Dose:** Usually **1–2 inhalations twice daily**, or as directed by the physician.
- Shake the inhaler well before each use.
- Exhale fully before placing the mouthpiece between the lips.
- Begin to inhale slowly and deeply while actuating the inhaler.
- Hold the breath for approximately **10 seconds**, or as comfortably possible, before exhaling.
- If another inhalation is prescribed, wait approximately **1 minute** before repeating.
- **Rinse the mouth with water and spit out after each dose** to reduce the risk of oral candidiasis and throat irritation.
- Use a **spacer device** when recommended.
- Do not stop regular treatment abruptly without medical advice.

Precautions:

- Use regularly as prescribed; it is **not intended for immediate relief of acute bronchospasm** unless the specific treatment plan allows formoterol-containing therapy for relief.
- Increasing use of a rescue inhaler or worsening symptoms may indicate inadequate disease control and requires medical review.
- Monitor patients for **oral candidiasis, hoarseness, tremor, palpitations, and changes in heart rate**.
- Long-term inhaled corticosteroid therapy may rarely affect adrenal function, bone health, eye health, or growth in children.
- Use cautiously in patients with **cardiovascular disease, hypertension, diabetes, hyperthyroidism, glaucoma, cataracts, osteoporosis, or a history of tuberculosis or other respiratory infections**.
- Ensure correct inhaler technique and regular cleaning of the mouthpiece.
- **Pregnancy and Lactation:** Use only when clearly indicated and under medical supervision.

Contraindications:

- Known hypersensitivity to **formoterol, fluticasone propionate, or any component of the formulation**.
- Not appropriate as the primary treatment for **acute severe asthma or status asthmaticus**.
- Acute bronchospasm requiring immediate intensive bronchodilator therapy should be treated with an appropriate rapid-relief medicine.

Adverse Reactions:

Common:

- Hoarseness

- Throat irritation
- Oral candidiasis (thrush)
- Headache
- Tremor
- Palpitations
- Muscle cramps

Less Common but Serious:

- Paradoxical bronchospasm
- Significant tachycardia or cardiac arrhythmia
- Hypokalemia
- Hyperglycemia
- Adrenal suppression with prolonged/high-dose corticosteroid exposure
- Increased susceptibility to certain infections
- Hypersensitivity reactions

Seek urgent medical attention if breathing suddenly worsens immediately after inhalation.

Drug Interactions:

- Other **β_2-agonists** may increase cardiovascular and metabolic adverse effects.
- **Non-selective beta-blockers** may reduce the bronchodilator effect and may precipitate bronchospasm.
- **MAO inhibitors and tricyclic antidepressants** may enhance cardiovascular effects of formoterol.
- **Diuretics** may increase the risk of hypokalemia.
- Other **corticosteroids** may increase the risk of systemic corticosteroid-related adverse effects.
- Strong **CYP3A4 inhibitors** may increase systemic exposure to fluticasone propionate.
- Certain medicines that prolong the **QT interval** may increase arrhythmia risk when combined with formoterol.

Storage:

- Store according to the manufacturer's specified conditions, generally below **25°C**.
- Protect from direct sunlight and excessive heat.
- Do not puncture, crush, or incinerate the canister.
- Keep the inhaler out of reach of children.
- Replace the protective cap after each use.
- Keep the mouthpiece clean.

Note:

This information is intended as a general prescribing guide. **Flutialfa F 250 Inhaler should be used under the supervision of a registered medical practitioner and with correct inhaler technique.** Regular use helps control airway inflammation and bronchoconstriction; it should not be discontinued without medical advice.

Important: The exact dosing schedule should follow the approved product label and the patient's asthma/COPD treatment plan. **Patients should have an appropriate rapid-relief bronchodilator available for acute symptoms unless their physician has specifically prescribed a formoterol-containing regimen for both maintenance and relief.**

Manufactured in India for:

Cafoli™
LIFECARE

Cafoli Lifecare Pvt. Ltd.
(An ISO 9001: 2015 Certified Co.)
Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113
Corporate Office :
Plot no. 307, Industrial Area Phase-2,
Panchkula, Haryana -134113
TM: Trademark Applied for