Information For the Use of
Fluticasone Furoate (100mcg) + Vilanterol (25mcg) Inhalation

Flutialfa VT 100 Inhalation

FLUTIALFA VT 100 INHALATION

Composition

Each delivered dose contains:

Vilanterol 25 mcg
Fluticasone Furoate 100 mcg
Excipients … q.s.

Pharmacological Action

Fluticasone Furoate is a potent inhaled corticosteroid (ICS) with anti-inflammatory activity. It acts locally in the respiratory tract to reduce airway inflammation, mucosal oedema and inflammatory responses associated with asthma.

Vilanterol is a long-acting selective β_2-adrenergic receptor agonist (LABA). It stimulates β_2-receptors in bronchial smooth muscle, resulting in increased intracellular cyclic AMP and prolonged relaxation of airway smooth muscle.

The combination provides complementary **anti-inflammatory and long-acting bronchodilator effects**, helping improve airway function and maintain respiratory symptom control.

Indications

Flutialfa VT 100 Inhalation is indicated for the **regular maintenance treatment of asthma** in patients for whom treatment with an inhaled corticosteroid and a long-acting β_2-agonist is appropriate.

It may be used in patients:

- Not adequately controlled with an inhaled corticosteroid and an as-needed short-acting β_2-agonist.
- Already adequately controlled on an inhaled corticosteroid and LABA.
- Requiring regular maintenance therapy for improved asthma control.

The 100/25 mcg strength may be selected according to the severity of asthma and the patient's response to treatment.

Important: Flutialfa VT 100 is **not intended for the relief of acute bronchospasm or sudden asthma attacks**. A suitable short-acting rescue bronchodilator should be available for acute symptoms.

Dosage and Administration

Adults:
The recommended dose is **one inhalation of Fluticasone Furoate 100 mcg + Vilanterol 25 mcg once daily**, administered by oral inhalation.

- Use at approximately the **same time each day**.
- Do not use more than **one inhalation in 24 hours**.
- Do not use an additional dose for sudden breathlessness.
- Use the prescribed short-acting rescue inhaler for acute symptoms.
- **Rinse the mouth with water and spit it out after inhalation** to reduce the risk of oral candidiasis.

Use the inhaler with the correct technique according to the instructions supplied with the particular inhaler device.

Precautions

- This product is **not a rescue medication** and should not be used for acute severe bronchospasm.
- LABA therapy should not be used as monotherapy for asthma; vilanterol is combined with an inhaled corticosteroid in this formulation.
- Monitor for oral or oropharyngeal candidiasis and dysphonia.
- Use cautiously in patients with cardiovascular disease, cardiac arrhythmias, hypertension or a history of QT-interval prolongation.
- Use cautiously in patients with hyperthyroidism, diabetes mellitus, seizure disorders or hypokalaemia.
- β_2-agonists may cause transient hypokalaemia, palpitations or changes in blood glucose.
- Prolonged corticosteroid exposure may be associated with systemic effects including adrenal suppression, decreased bone mineral density, cataract and glaucoma.
- Patients with significant hepatic impairment may have increased systemic exposure to fluticasone furoate and should be monitored appropriately.

Contraindications

Flutialfa VT 100 Inhalation is contraindicated in patients with:

- Hypersensitivity to **fluticasone furoate, vilanterol** or any component of the formulation.
- Severe hypersensitivity to milk proteins, where applicable to a lactose-containing dry-powder inhaler.
- Primary treatment of status asthmaticus or other acute episodes of asthma requiring intensive treatment.

Adverse Reactions

Common/possible:

- Headache
- Nasopharyngitis
- Upper respiratory tract infection
- Cough
- Hoarseness/dysphonia
- Oral or oropharyngeal candidiasis
- Tremor
- Palpitations
- Muscle cramps

Other/serious possible reactions:

- Paradoxical bronchospasm
- Tachycardia or cardiac arrhythmias
- Hypokalaemia
- Hyperglycaemia
- Hypersensitivity reactions
- Adrenal suppression with excessive systemic corticosteroid exposure
- Cataract or glaucoma with prolonged corticosteroid exposure

Drug Interactions

Strong CYP3A4 inhibitors:
Medicines such as ketoconazole, ritonavir and cobicistat may increase systemic exposure to fluticasone furoate and vilanterol. Concomitant use requires caution.

β-blockers:
May antagonise the bronchodilator effect of vilanterol and may cause bronchospasm. Use together only when clinically necessary and under medical supervision.

MAO inhibitors and tricyclic antidepressants:
May potentiate cardiovascular effects associated with β_2-agonists.

Non-potassium-sparing diuretics:
May increase the risk of β_2-agonist-associated hypokalaemia and ECG changes.

Other medicines capable of prolonging the QT interval should be used cautiously.

Pregnancy and Lactation

Use during pregnancy should be considered when the expected benefit to the mother justifies potential risk to the fetus.

During breastfeeding, treatment should be continued only following appropriate medical assessment of maternal benefit and potential risk to the infant.

Storage

Store the inhaler according to the **approved device-specific storage conditions**.

Keep in the original packaging and protect from moisture, heat and direct sunlight.

Keep out of reach of children.

Do not use after the expiry date or if the inhaler/device is damaged.

Note

For use under medical supervision.

Flutialfa VT 100 Inhalation is a **maintenance controller medication and not a rescue inhaler**. It should be used regularly as prescribed and should not be administered more frequently than the recommended once-daily dose. Correct inhaler technique and regular clinical assessment are important for optimal asthma control.

Manufactured in India for:

Cafoli™
LIFECARE

Cafoli Lifecare Pvt. Ltd.
(An ISO 9001: 2015 Certified Co.)
Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113
Corporate Office :
Plot no. 307, Industrial Area Phase-2,
Panchkula, Haryana -134113
TM: Trademark Applied for