Information For the Use of
Fosfomycin (3gm) Sachet

Fosfohub Sachet

Fosfohub Sachet

Composition:

Each sachet contains:

- **Fosfomycin** – 3 g

Pharmacological Action:

Fosfomycin is a **bactericidal antibiotic** that acts at an early stage of bacterial cell-wall synthesis.

- **Cell-Wall Inhibition:** Fosfomycin inhibits the enzyme MurA, blocking an essential step in peptidoglycan synthesis.
- **Bactericidal Action:** Disruption of bacterial cell-wall formation leads to bacterial cell death.
- **Urinary Tract Activity:** After oral administration, fosfomycin achieves high concentrations in urine, making it particularly useful against susceptible urinary pathogens.
- **Broad Antibacterial Activity:** Active against several Gram-negative and Gram-positive urinary pathogens, including susceptible *E. coli*.

Indications:

Fosfohub Sachet is indicated for the treatment of:

- **Acute Uncomplicated Cystitis in Adult Women**
- **Lower Urinary Tract Infection (UTI)** caused by susceptible organisms
- Other susceptible urinary infections when specifically prescribed by the physician

---

Dosage and Administration:

- **Recommended Dose: One sachet (3 g) as a single dose** for acute uncomplicated cystitis in adult women, or as directed by the physician.
- Empty the contents of the sachet into a glass of water and stir until completely dissolved.
- Drink immediately after preparation.
- Preferably take on an **empty stomach**, approximately 2–3 hours before or after a meal, or at bedtime after emptying the bladder.
- Do not take repeated doses unless specifically prescribed by the physician.

Precautions:

- Fosfomycin should be used only for infections where it is appropriate and susceptibility is expected.
- It is **not generally appropriate for pyelonephritis or complicated systemic urinary infections** because oral fosfomycin does not achieve adequate renal tissue concentrations.
- Patients with fever, chills, flank/back pain, vomiting, or systemic symptoms should seek medical evaluation for possible upper urinary tract infection.
- Use cautiously in patients with significant renal impairment.
- Complete the prescribed treatment and avoid unnecessary antibiotic use to reduce antimicrobial resistance.
- **Pregnancy and Lactation:** Fosfomycin may be used when clinically indicated; treatment should be guided by a healthcare professional.

Contraindications:

- Known hypersensitivity to **fosfomycin** or any component of the formulation.
- Severe renal impairment where oral fosfomycin use is not appropriate.
- Hereditary **fructose intolerance or other relevant excipient-related disorders**, if the particular sachet contains sucrose or fructose-containing excipients.

Adverse Reactions:

Common:

- Diarrhea
- Nausea
- Abdominal pain
- Headache
- Indigestion
- Vaginitis

Less Common but Serious:

- Severe or persistent diarrhea associated with **Clostridioides difficile**
- Severe hypersensitivity reactions
- Angioedema
- Anaphylaxis
- Significant skin reactions

Seek immediate medical attention for breathing difficulty, facial swelling, severe rash, or persistent severe diarrhea.

Drug Interactions:

- **Metoclopramide** may reduce fosfomycin concentrations and can decrease its urinary exposure; concomitant use should generally be avoided unless specifically advised.
- Other medicines that significantly alter gastrointestinal motility may potentially affect absorption.
- Inform the physician about all prescription medicines, OTC medicines, and supplements being used.

Storage:

- Store in a cool, dry place below **25°C**.
- Protect from moisture and direct sunlight.
- Keep the sachet sealed until use.
- Keep out of reach of children.

Note:

This information is intended as a general prescribing guide. Fosfohub Sachet should be used under the supervision of a registered medical practitioner. **Fosfomycin 3 g oral sachet is primarily used as a single-dose treatment for acute uncomplicated cystitis in adult women.** It should not be routinely used for kidney infections or complicated UTIs without specialist advice.

Important: Antibiotic selection should be based on the suspected pathogen, local resistance patterns, and clinical assessment.

Manufactured in India for:

Cafoli™ LIFECARE

Cafoli Lifecare Pvt. Ltd.
(An ISO 9001: 2015 Certified Co.)
Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113
Corporate Office :
Plot no. 307, Industrial Area Phase-2,
Panchkula, Haryana -134113
TM: Trademark Applied for