Information For the Use of
Gatifloxacin (0.3% w/v) + Prednisolone (1% w/v) Eye Drops

Gatitrip P Eye Drop

Gatitrip P Eye Drops

Composition:

Each mL contains:

- **Gatifloxacin** – 0.3% w/v
- **Prednisolone** – 1% w/v

Pharmacological Action:

Gatitrip P Eye Drops combines a **fluoroquinolone antibacterial** with a **corticosteroid** to provide antibacterial and anti-inflammatory activity in the eye.

- **Antibacterial Action:** Gatifloxacin inhibits bacterial DNA gyrase and topoisomerase IV, interfering with bacterial DNA replication and causing bacterial cell death.
- **Anti-inflammatory Action:** Prednisolone suppresses the release and activity of inflammatory mediators, reducing ocular inflammation.
- **Combined Action:** Provides antimicrobial coverage while controlling associated inflammatory symptoms in appropriate bacterial inflammatory eye conditions.

Indications:

Gatitrip P Eye Drops is indicated for the treatment of **steroid-responsive ocular inflammation when a bacterial infection or risk of bacterial infection is present**, including:

- **Bacterial Conjunctivitis with Inflammation**
- **Blepharoconjunctivitis**
- **Post-operative Ocular Inflammation with Bacterial Infection/Risk**
- Other susceptible bacterial ocular infections associated with inflammation, as prescribed by an ophthalmologist

---

Dosage and Administration:

- **Recommended Dose:** Instill the prescribed number of drops into the affected eye(s) as directed by the ophthalmologist.
- Wash hands thoroughly before use.
- Shake well if the formulation requires shaking.
- Avoid touching the dropper tip to the eye, eyelids, or surrounding surfaces.

- Close the eye gently after instillation and avoid excessive blinking.
- If other ophthalmic medicines are being used, maintain an appropriate interval between applications.
- Do not use for longer than prescribed because prolonged topical corticosteroid therapy may increase ocular complications.

Precautions:

- For **ophthalmic use only**.
- Use under ophthalmic supervision, particularly when treatment extends beyond a short course.
- Prolonged corticosteroid use may increase **intraocular pressure** and the risk of glaucoma or cataract.
- Corticosteroids may delay corneal wound healing and can mask or worsen certain ocular infections.
- Use cautiously in patients with a history of **glaucoma, ocular herpes infection, or other viral/fungal eye infections**.
- Contact lens wear should be avoided during active bacterial conjunctivitis unless specifically advised by the ophthalmologist.
- **Pregnancy and Lactation:** Use only when clearly indicated and under medical supervision.

Contraindications:

- Known hypersensitivity to **gatifloxacin, prednisolone**, other quinolones, corticosteroids, or any component of the formulation.
- **Herpes simplex keratitis** or other viral infections of the eye where corticosteroid use is contraindicated.
- Fungal diseases of ocular structures.
- Untreated mycobacterial infections of the eye.
- Acute untreated purulent ocular infections where corticosteroid therapy is inappropriate.

Adverse Reactions:

Common:

- Temporary burning or stinging
- Eye irritation
- Blurred vision immediately after instillation
- Mild redness or discomfort

Less Common but Serious:

- Increased intraocular pressure

- Glaucoma
- Cataract formation with prolonged use
- Delayed corneal wound healing
- Secondary ocular infection
- Corneal or scleral thinning in susceptible patients
- Severe hypersensitivity reactions

Patients should seek immediate ophthalmic evaluation if eye pain, worsening redness, reduced vision, marked swelling, or severe irritation develops.

Drug Interactions:

- Concurrent use with other **ophthalmic corticosteroids** may increase the risk of steroid-related adverse effects.
- Other topical ophthalmic medicines should preferably be administered with an appropriate interval between applications.
- Systemic absorption of ophthalmic gatifloxacin and prednisolone is limited, but the treating physician should be informed about all medicines being used.

Storage:

- Store in a cool, dry place below **25°C**.
- Protect from direct sunlight and excessive heat.
- Keep the bottle tightly closed when not in use.
- Do not touch the dropper tip to any surface.
- Keep out of reach of children.

Note:

This information is intended as a general prescribing guide. Gatitrip P Eye Drops should be used only under the supervision of a qualified ophthalmologist. Because it contains a corticosteroid, prolonged or inappropriate use may increase intraocular pressure and may mask or worsen certain ocular infections.

Because Gatitrip P contains a corticosteroid, safe use depends on limiting treatment duration and monitoring for steroid-related complications such as increased intraocular pressure and worsening infection.

Manufactured in India for:

Cafoli Lifecare Pvt. Ltd.
(An ISO 9001: 2015 Certified Co.)
Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113
Corporate Office :
Plot no. 307, Industrial Area Phase-2,
Panchkula, Haryana -134113
TM: Trademark Applied for