Information For the Use of
Glimepiride (2mg) + Metformin (850mg) SR Bilayered Tablet

Gimtype M2/850 Tablet

GIMTYPE M2/850 TABLETS

Composition

Each film-coated bilayered tablet contains:

Glimepiride I.P. 2 mg
Metformin Hydrochloride I.P. 850 mg
(as Sustained Release)
Excipients … q.s.

Pharmacological Action

Glimepiride is a second-generation sulfonylurea. It lowers blood glucose primarily by stimulating pancreatic β-cells to release insulin. It also improves glucose utilization and contributes to reduction of postprandial and fasting blood glucose.

Metformin Hydrochloride is a biguanide antihyperglycaemic agent. It primarily reduces hepatic glucose production, decreases intestinal glucose absorption to a lesser extent and improves peripheral insulin sensitivity, thereby increasing glucose uptake and utilization.

The combination provides complementary antihyperglycaemic action through **increased insulin secretion and improved insulin sensitivity/reduced hepatic glucose production**.

Indications

Gimtype M2/850 Tablets are indicated as an adjunct to **diet and exercise** for the management of **Type 2 Diabetes Mellitus** in adults when adequate glycaemic control is not achieved with appropriate lifestyle measures and/or when combination therapy with glimepiride and metformin is considered appropriate.

The combination should be used according to the patient's individual glycaemic response and the approved product-specific prescribing information. CDSCO records include glimepiride + sustained-release metformin combinations for Type 2 diabetes when diet, exercise and single-agent therapy do not provide adequate glycaemic control.

Dosage and Administration

Adults:

Dosage should be individualized according to blood glucose levels, HbA1c, renal function, tolerability and previous antidiabetic therapy.

For patients being transferred from separate glimepiride and metformin therapy, the combination strength should correspond as closely as possible to the established individual doses.

Administration:

- Take the tablet **with meals**, preferably with the main meal, or as directed by the physician.
- Swallow the tablet whole with sufficient water.
- **Do not crush, chew or break** the sustained-release bilayered tablet.
- Regular monitoring of fasting blood glucose and HbA1c is recommended.

Dose adjustment may be necessary in elderly patients and patients with renal impairment.

Precautions

- **Hypoglycaemia** may occur, particularly because of the glimepiride component.
- Patients should be educated about recognizing symptoms of hypoglycaemia such as sweating, tremor, hunger, dizziness, palpitations, weakness and confusion.
- Renal function should be assessed before initiation and monitored periodically during treatment.
- Metformin should not be used in patients with severe renal impairment.
- Monitor hepatic function in patients with significant liver disease.
- Excessive alcohol intake should be avoided because it may increase the risk of hypoglycaemia and metformin-associated lactic acidosis.
- Temporarily discontinue metformin-containing therapy when clinically appropriate around certain **iodinated contrast** procedures, with renal function reassessed before restarting.
- Withhold treatment during significant conditions associated with hypoxia, dehydration, severe infection or acute illness when clinically indicated.
- Elderly or debilitated patients require careful dose selection because of increased susceptibility to hypoglycaemia and renal impairment.
- Maintain regular dietary intake while taking glimepiride.

Contraindications

Gimtype M2/850 Tablets are contraindicated in:

- Known hypersensitivity to **glimepiride, metformin or any component** of the formulation.
- **Severe renal impairment** or significantly reduced kidney function where metformin is contraindicated.
- Acute or chronic metabolic acidosis, including **diabetic ketoacidosis**.
- Severe hepatic impairment where use is inappropriate.
- Conditions associated with tissue hypoxia or severe circulatory/respiratory compromise where metformin-associated lactic acidosis risk is increased.

- Type 1 diabetes mellitus.

Adverse Reactions

Common/Possible:

- Hypoglycaemia
- Nausea
- Vomiting
- Diarrhoea
- Abdominal discomfort
- Dyspepsia
- Loss of appetite
- Metallic taste

Other possible reactions:

- Headache
- Dizziness
- Weakness
- Skin rash or hypersensitivity reactions

Rare/Serious:

- Severe or prolonged hypoglycaemia
- Lactic acidosis associated with metformin, particularly in high-risk patients
- Hepatic or haematological abnormalities
- Vitamin B12 deficiency with prolonged metformin treatment

Drug Interactions

Medicines increasing hypoglycaemic effect:
Insulin and other antidiabetic medicines, some antibiotics and other medicines may increase the risk of hypoglycaemia.

Medicines reducing glycaemic control:
Certain corticosteroids, sympathomimetic agents, thiazide-type diuretics and other medicines may increase blood glucose and reduce the glucose-lowering effect.

Beta-blockers:
May mask some warning symptoms of hypoglycaemia.

Alcohol:
Excessive alcohol consumption may increase the risk of hypoglycaemia and metformin-associated lactic acidosis.

Iodinated contrast media:
Metformin may need to be temporarily discontinued around iodinated contrast procedures depending on renal function and the clinical situation.

Cationic drugs:
Medicines eliminated through renal tubular secretion may potentially interact with metformin and should be reviewed by the physician.

Patients should inform their healthcare professional about all prescription, OTC and herbal medicines being taken.

Pregnancy and Lactation

Gimtype M2/850 Tablets should generally **not be used during pregnancy unless specifically prescribed by the treating physician**. Insulin is generally preferred when adequate glycaemic control is required during pregnancy.

Glimepiride and metformin can pass into breast milk to varying degrees. The decision to use this combination during lactation should be made by the physician after considering the benefits and potential risks.

Storage

Store in a **cool, dry place**, protected from excessive heat, moisture and direct sunlight.

Keep the tablets in their original packaging and **out of reach of children**.

Do not use after the expiry date.

Note

For use under medical supervision.

Gimtype M2/850 Tablets should be used as an adjunct to appropriate **diet and exercise**. Blood glucose and HbA1c should be monitored regularly, with particular attention to renal function and symptoms of hypoglycaemia.

Important: The final printed literature should follow the **approved product-specific Indian prescribing information**, particularly for exact dosage, renal cut-offs, contraindications and approved indications.

Manufactured in India for:

Cafoli Lifecare Pvt. Ltd.
(An ISO 9001: 2015 Certified Co.)
Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113
Corporate Office :
Plot no. 307, Industrial Area Phase-2,
Panchkula, Haryana -134113
TM: Trademark Applied for