Information For the Use of
Gliclazide (80mg) + Metformin (500mg) ER Bilayered Tablet

Glicaturn M 80 Tablet

GLICATURN M 80 TABLET

Gliclazide (80 mg) + Metformin (500 mg) ER Bilayered Tablet

1. Composition

Each bilayered tablet contains:
Gliclazide 80 mg + Metformin Hydrochloride 500 mg ER
Excipients q.s.

2. Pharmacological Action

Gliclazide is a sulfonylurea antidiabetic agent. It stimulates pancreatic β-cells to release insulin and thereby helps reduce blood glucose levels.

Metformin is a biguanide that primarily decreases hepatic glucose production and improves peripheral insulin sensitivity, thereby enhancing glucose utilization.

The combination provides complementary mechanisms for improving **glycaemic control in patients with Type 2 diabetes mellitus**.

3. Indications

Glicaturn M 80 Tablet is indicated as an adjunct to diet and exercise for the treatment of **Type 2 diabetes mellitus** when adequate glycaemic control is not achieved with lifestyle measures or single-agent therapy, as appropriate.

It may help:

- Reduce fasting and postprandial blood glucose.
- Improve overall glycaemic control.
- Reduce HbA1c in appropriately selected patients.

4. Dosage and Administration

The dose should be individualized according to the patient's blood glucose levels, renal function, tolerability and previous antidiabetic therapy.

Usual administration: One tablet once daily or as directed by the physician. Depending on the patient's requirements, the dose may be adjusted under medical supervision.

The tablet should be taken **with or immediately after a meal** and swallowed whole with water. The ER portion should not be crushed or chewed.

Patients should maintain their prescribed diet and exercise programme.

5. Precautions

- Monitor blood glucose regularly, particularly when therapy is initiated or adjusted.
- **Hypoglycaemia** may occur, particularly with gliclazide, if meals are skipped, caloric intake is inadequate, alcohol is consumed, or other glucose-lowering medicines are used.
- Assess renal function before starting treatment and periodically thereafter.
- Metformin should be used cautiously or discontinued when clinically appropriate in conditions associated with significant renal dysfunction or tissue hypoxia.
- Avoid excessive alcohol consumption because it may increase the risk of hypoglycaemia and metformin-associated lactic acidosis.
- Temporarily withhold metformin-containing therapy when clinically indicated around certain iodinated contrast procedures, according to renal function and current prescribing recommendations.
- Monitor hepatic function where clinically appropriate.
- Inform the physician about any planned surgery or prolonged fasting.

6. Contraindications

The combination is contraindicated in patients with:

- Known hypersensitivity to **gliclazide, metformin, sulfonylureas, sulfonamides**, or any component of the formulation.
- **Type 1 diabetes mellitus** or diabetic ketoacidosis.
- Severe renal impairment or acute conditions associated with impaired renal function, where metformin is contraindicated.
- Acute or chronic metabolic acidosis, including lactic acidosis.
- Severe conditions associated with tissue hypoxia, where metformin therapy is contraindicated according to approved prescribing information.
- Severe hepatic impairment where use is contraindicated.

7. Adverse Reactions

Possible adverse reactions include:

Gliclazide-related:

- Hypoglycaemia
- Headache
- Dizziness
- Gastrointestinal discomfort
- Weight gain

Metformin-related:

- Nausea and vomiting
- Diarrhoea
- Abdominal discomfort
- Loss of appetite
- Metallic taste

Long-term metformin therapy may be associated with **vitamin B12 deficiency**.

Rare but serious: **lactic acidosis**, particularly in patients with significant risk factors or contraindications to metformin.

8. Drug Interactions

Important interactions may occur with:

- Other antidiabetic medicines, increasing the risk of **hypoglycaemia**.
- Alcohol, which may increase the risk of hypoglycaemia and lactic acidosis.
- Certain medicines such as corticosteroids, sympathomimetics and some diuretics, which may increase blood glucose.
- ACE inhibitors and other medicines that can alter glucose control may require monitoring.
- Cationic medicines eliminated by renal tubular secretion may increase metformin exposure.
- Iodinated contrast media may increase the risk of metformin-associated complications in susceptible patients.

The glucose-lowering effect of gliclazide may be increased by certain antibiotics, anticoagulants and other medicines; therefore, appropriate monitoring is advised.

9. Storage

Store below **30°C**, protected from light and moisture.

Keep out of reach of children.

10. Note / Important

Glicaturn M 80 Tablet is intended for the management of **Type 2 diabetes mellitus** along with appropriate diet and exercise.

Hypoglycaemia is an important potential adverse effect of gliclazide. Patients should be educated about recognition and management of low blood glucose.

Renal function should be assessed regularly during metformin therapy. The dosage and use of the combination should be individualized according to the patient's clinical response and the approved prescribing information.

For use under medical supervision.

Keep out of reach of children.

Manufactured in India for:

Cafoli Lifecare Pvt. Ltd.
(An ISO 9001: 2015 Certified Co.)
Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113
Corporate Office :
Plot no. 307, Industrial Area Phase-2,
Panchkula, Haryana -134113
TM: Trademark Applied for