Information For the Use of
Granisetron (1mg) MD Tablets

Granityle MD Tablet

GRANITYLE MD TABLET

Granisetron 1 mg MD

Composition

Each orally disintegrating tablet contains:

- **Granisetron – 1 mg**

Pharmacological Action

Granisetron is a potent and selective **5-HT_3 (serotonin) receptor antagonist** with antiemetic activity. It prevents nausea and vomiting by blocking serotonin-mediated stimulation of 5-HT_3 receptors in the gastrointestinal tract and central nervous system.

It functions by:

- **5-HT_3 Receptor Blockade:** Selectively blocks serotonin at 5-HT_3 receptors involved in the vomiting reflex.
- **Antiemetic Action:** Helps prevent and control nausea and vomiting associated with cancer treatment.
- **Chemotherapy Support:** Reduces chemotherapy-induced nausea and vomiting.
- **Radiotherapy Support:** Helps prevent nausea and vomiting associated with radiation therapy.

Indications

Granityle MD Tablets are indicated for the prevention and management of:

- **Chemotherapy-induced nausea and vomiting (CINV)**
- **Radiation-induced nausea and vomiting**
- Nausea and vomiting associated with highly emetogenic cancer chemotherapy, including high-dose cisplatin
- Other conditions associated with nausea and vomiting, as prescribed by the physician.

Dosage and Administration

- **Recommended Dose:** For adults receiving emetogenic chemotherapy, the usual oral regimen is **2 mg once daily or 1 mg twice daily**, as directed by the physician.
- **Timing:** The first dose is generally administered up to **1 hour before chemotherapy**. With the 1 mg twice-daily regimen, the second dose is taken approximately 12 hours after the first.
- **Administration:** Place the MD tablet on the tongue and allow it to disintegrate before swallowing. Water may be taken if required.

- **With or Without Food:** Granisetron may be taken with or without food.
- **Duration of Use:** Use according to the chemotherapy/radiotherapy schedule and physician's advice.

Precautions

- **Hypersensitivity:** Use cautiously in patients with a history of hypersensitivity to granisetron or other 5-HT_3 receptor antagonists.
- **Cardiac Conditions:** Granisetron may affect cardiac repolarization, including QT interval; use cautiously in patients with cardiac disease or risk factors for QT prolongation.
- **Electrolyte Abnormalities:** Correct significant electrolyte disturbances before treatment where appropriate.
- **Gastrointestinal Conditions:** Granisetron may reduce intestinal motility; patients with signs of intestinal obstruction should be monitored carefully.
- **Serotonin Syndrome:** Monitor patients receiving granisetron with serotonergic medicines for symptoms suggestive of serotonin syndrome.
- **Pregnancy & Lactation:** Use during pregnancy or breastfeeding only when considered necessary by the treating physician.

Contraindications

- Known hypersensitivity to **granisetron** or any component of the formulation.
- Patients with a previous serious hypersensitivity reaction to granisetron or other 5-HT_3 receptor antagonists should not receive the medicine unless specifically evaluated by a physician.

Adverse Reactions

Common:

- Headache
- Constipation
- Weakness or fatigue
- Dizziness
- Abdominal discomfort

Rare but Serious:

- Hypersensitivity reactions
- QT prolongation and cardiac rhythm disturbances
- Serotonin syndrome, particularly with concomitant serotonergic medicines
- Severe allergic reactions

Patients should seek medical attention for severe allergic symptoms, abnormal heartbeat, severe dizziness, agitation, muscle rigidity or other unusual symptoms.

Drug Interactions

- **Serotonergic Medicines:** Concomitant use with SSRIs, SNRIs, MAO inhibitors, mirtazapine, tramadol and other serotonergic medicines may increase the risk of serotonin syndrome.
- **QT-Prolonging Medicines:** Caution is advised with medicines that may prolong the QT interval or cause cardiac rhythm disturbances.
- **Chemotherapy Medicines:** Inform the treating physician about all anticancer medicines being administered.
- Patients should inform their physician about all prescription, OTC and herbal medicines being taken before starting treatment.

Storage

- Store in a **cool, dry place**, protected from moisture and excessive heat.
- Protect the tablets from light and retain them in the original packaging until use.
- Keep out of reach of children.
- Follow the storage conditions specified on the product packaging.

Note

This information is intended as a **professional pharmaceutical-literature guideline**. **Granityle MD should be used under the supervision of a qualified healthcare professional.** The exact dosage and duration should be determined according to the indication, chemotherapy/radiotherapy regimen and individual patient requirements. The Indian Pharmacopoeia Commission recognizes granisetron tablets in **1 mg and 2 mg strengths** and classifies granisetron hydrochloride as a 5-HT_3 receptor antagonist used in the treatment of nausea and vomiting.

Manufactured in India for:

Cafoli™
LIFECARE

Cafoli Lifecare Pvt. Ltd.
(An ISO 9001: 2015 Certified Co.)
Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113
Corporate Office :
Plot no. 307, Industrial Area Phase-2,
Panchkula, Haryana -134113
TM: Trademark Applied for