Haloside S Ointment

HALOSIDE S OINTMENT

Composition

Each gram of ointment contains:

Halobetasol Propionate I.P. 0.05% w/w
Salicylic Acid I.P. 3% w/w
Ointment base … q.s.

Pharmacological Action

Halobetasol Propionate is a super-potent topical corticosteroid. It produces anti-inflammatory, antipruritic and vasoconstrictive effects by reducing the release and activity of inflammatory mediators in the skin.

Salicylic Acid is a topical keratolytic agent. It softens and removes excessive scales and keratin, helping to reduce hyperkeratosis and allowing better penetration of the corticosteroid into thickened lesions.

The combination therefore provides **anti-inflammatory, antipruritic and keratolytic action**, particularly useful in inflammatory dermatoses associated with scaling and hyperkeratosis.

Indications

Haloside S Ointment may be used, under medical supervision, for short-term topical treatment of corticosteroid-responsive inflammatory and hyperkeratotic skin conditions such as:

- Psoriasis with thick scaling
- Lichenified eczema/dermatitis
- Hyperkeratotic eczema
- Other steroid-responsive dermatoses associated with excessive scaling

Use should be limited to conditions for which a super-potent topical corticosteroid is appropriate.

Dosage and Administration

Adults:
Apply a **thin film** of Haloside S Ointment to the affected area **once or twice daily**, or as directed by the physician.

Gently rub into the affected skin until absorbed.

Duration:
Because Halobetasol Propionate is a **super-high-potency corticosteroid**, treatment should generally be limited to the shortest duration necessary to achieve control. Prolonged continuous use should be avoided unless specifically directed by the physician.

Do not apply more frequently or over larger areas than recommended.

Precautions

- For **external use only**.
- Avoid contact with the **eyes, mouth and mucous membranes**.
- Do not apply to the face, groin or axillae unless specifically directed by a physician.
- Avoid application to large areas of the body or under occlusive dressings unless prescribed.
- Prolonged or excessive use may cause **skin atrophy, striae and other local corticosteroid effects**.
- Salicylic acid may cause irritation, particularly when applied to sensitive or damaged skin.
- Use cautiously on broken, infected or severely inflamed skin.
- If irritation, excessive dryness or sensitization develops, treatment should be reassessed.
- Children may be more susceptible to systemic corticosteroid effects; use only under appropriate medical supervision.

Contraindications

Haloside S Ointment is contraindicated in:

- Known hypersensitivity to **Halobetasol Propionate, Salicylic Acid or any component of the formulation**.
- Untreated primary bacterial, fungal or viral skin infections.
- Tuberculosis or other specific infections of the skin where topical corticosteroid treatment is inappropriate.
- Rosacea or perioral dermatitis where corticosteroid therapy is contraindicated.

It should not be used for conditions not responsive to topical corticosteroids.

Adverse Reactions

Common/Local:

- Burning or stinging
- Itching
- Irritation
- Dryness
- Redness

With prolonged or inappropriate use:

- Skin thinning (atrophy)
- Striae
- Telangiectasia
- Folliculitis
- Acneiform eruptions
- Hypopigmentation
- Contact dermatitis

Rare/Systemic:

With excessive application, prolonged use or application over large areas, systemic corticosteroid effects including **hypothalamic-pituitary-adrenal (HPA) axis suppression** may occur.

Drug Interactions

Clinically significant interactions are uncommon with appropriate topical use.

However, concomitant use of other **topical corticosteroids, keratolytic agents or irritating dermatological preparations** on the same area may increase local irritation or corticosteroid exposure.

Patients should inform their physician about other topical and systemic medicines being used.

Pregnancy and Lactation

Use during pregnancy should be considered only when the expected benefit justifies the potential risk.

Because Halobetasol Propionate is a very potent corticosteroid, prolonged or extensive use during pregnancy should be avoided unless specifically prescribed.

During lactation, use only on medical advice. **Do not apply to the breast or nipple immediately before breastfeeding.** If applied to the breast, the area should be thoroughly cleansed before feeding.

Storage

Store in a **cool, dry place**, protected from excessive heat and direct sunlight.

Keep the container tightly closed and **out of reach of children**.

Do not use after the expiry date.

Note

For external use only. For use under medical supervision.

Apply only to the affected skin and use for the shortest duration necessary. Avoid prolonged, extensive or occlusive use unless specifically prescribed.

Important: Product-specific approved prescribing information should take precedence for the final printed literature, particularly regarding duration of treatment, age restrictions and maximum quantity/application area.

Manufactured in India for:

Cafoli Lifecare Pvt. Ltd.
(An ISO 9001: 2015 Certified Co.)
Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113
Corporate Office :
Plot no. 307, Industrial Area Phase-2,
Panchkula, Haryana -134113
TM: Trademark Applied for