Information For the Use of
Ademetionine (200mg) Tablets

Cafosam 200 Tablet

CAFOSAM 200 TABLET

Ademetionine (200 mg)

1. Composition

Each enteric-coated tablet contains:
Ademetionine 200 mg
Excipients q.s.

2. Pharmacological Action

Ademetionine, also known as **S-Adenosyl-L-Methionine (SAMe)**, is a naturally occurring compound involved in transmethylation, transsulfuration and aminopropylation reactions.

It participates in several biochemical processes in the liver and may support **hepatic cellular function and bile flow**. Ademetionine is used therapeutically in selected liver disorders associated with cholestasis.

3. Indications

Ademetionine may be indicated in the management of selected hepatic and cholestatic conditions, including:

- **Intrahepatic cholestasis**.
- Cholestasis associated with chronic liver disorders.
- Cholestasis associated with pregnancy, where clinically appropriate.
- Other liver conditions where ademetionine is specifically prescribed.

4. Dosage and Administration

The dosage should be individualized according to the indication and clinical response.

Adults: Oral doses commonly range from **400–1600 mg daily**, divided into one or more doses, as directed by the physician.

The tablet should be swallowed **whole**, preferably between meals. Do not crush or chew enteric-coated tablets.

The duration of treatment depends on the underlying condition and response to therapy.

5. Precautions

- Use cautiously in patients with **renal impairment**, as clinical experience may be limited.
- Patients with bipolar disorder should use ademetionine only under medical supervision, as mood elevation or mania may occur.
- Patients receiving antidepressants or other serotonergic medicines should be monitored for possible interactions.
- Monitor patients for nausea, gastrointestinal discomfort or other adverse effects during treatment.
- Ademetionine may interfere with certain laboratory measurements in some circumstances; inform the healthcare professional about its use.
- Use during pregnancy and lactation should be based on an appropriate benefit-risk assessment.

6. Contraindications

Ademetionine is contraindicated in patients with:

- Known hypersensitivity to **ademetionine or any component of the formulation**.
- Conditions in which use is specifically contraindicated according to the approved product information.

7. Adverse Reactions

Possible adverse reactions include:

- Nausea
- Abdominal discomfort
- Diarrhoea
- Headache
- Dizziness
- Insomnia
- Anxiety

Rarely, **hypersensitivity reactions or mood elevation/mania** may occur, particularly in susceptible patients.

8. Drug Interactions

Particular caution is advised with:

- **Selective serotonin reuptake inhibitors (SSRIs)**.
- Tricyclic antidepressants.
- Other medicines or supplements that increase serotonergic activity.

Concomitant use may rarely increase the risk of **serotonin syndrome**. Patients receiving antidepressant therapy should be monitored appropriately.

Inform the physician about all prescription, OTC and herbal medicines being taken.

9. Storage

Store below **30°C**, protected from light and moisture.

Keep the container tightly closed.

Keep out of reach of children.

10. Note / Important

Ademetionine is used as an adjunct in selected **cholestatic and liver-related disorders**. The dose and duration should be individualized according to the underlying condition and clinical response.

Treatment should not be used as a substitute for management of the underlying cause of liver disease.

For use under medical supervision.

Keep out of reach of children.

Manufactured in India for:

Cafoli™
LIFECARE

Cafoli Lifecare Pvt. Ltd.
(An ISO 9001: 2015 Certified Co.)
Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113
Corporate Office :
Plot no. 307, Industrial Area Phase-2,
Panchkula, Haryana -134113
TM: Trademark Applied for