Information For the Use of
Hyoscine Butylbromide 20 mg injection

Butyhype Injection

BUTYHYPE INJECTION

Composition

Each mL of injection contains:

Hyoscine Butylbromide 20 mg
Water for Injection … q.s.

Pharmacological Action

Hyoscine Butylbromide is a quaternary ammonium antimuscarinic agent with antispasmodic activity.

It acts predominantly on muscarinic receptors in the smooth muscle of the gastrointestinal and genitourinary tracts, reducing abnormal or painful smooth-muscle contractions.

Because of its quaternary structure, systemic penetration into the central nervous system is limited. It therefore produces primarily **peripheral anticholinergic and antispasmodic effects**.

Indications

Butyhype Injection is indicated for the short-term relief of **acute painful smooth-muscle spasms**, including:

- Gastrointestinal tract spasm
- Biliary tract spasm
- Renal/ureteric colic
- Genitourinary tract spasm
- Other acute visceral smooth-muscle spasms where an antispasmodic is clinically appropriate

It may also be used in selected diagnostic or therapeutic procedures where reduction of smooth-muscle spasm is required.

Dosage and Administration

Adults and adolescents over 12 years:

The usual dose is:

20–40 mg by slow intravenous injection or intramuscular injection, as directed by the physician.

The dose may be repeated as clinically required, subject to the maximum recommended daily dose in the approved prescribing information.

Maximum daily dose:
Do not exceed **100 mg/day** unless specifically directed according to the applicable approved product information.

Administration:

- May be administered by **slow IV injection or IM injection**.
- IV administration should be given slowly because rapid injection may cause significant adverse cardiovascular effects.
- The patient should preferably be observed after parenteral administration.
- Use only when parenteral treatment is clinically indicated.

Precautions

- Use cautiously in patients with **tachycardia or cardiac disease**.
- Monitor patients receiving IV administration for hypotension, tachycardia or other cardiovascular effects.
- Use cautiously in patients with glaucoma, particularly **angle-closure glaucoma**.
- Use with caution in patients with suspected intestinal obstruction or urinary retention.
- Patients with prostatic enlargement may have an increased risk of urinary retention.
- Use cautiously in patients with significant hepatic or renal impairment.
- Parenteral anticholinergic therapy should not be continued without investigating the underlying cause of persistent or severe abdominal pain.
- If severe abdominal pain persists or is accompanied by fever, vomiting, blood in stool, fainting or marked tenderness, urgent medical evaluation is required.
- Patients receiving IV treatment should be monitored for **anaphylactic or severe hypersensitivity reactions**.

Contraindications

Butyhype Injection is contraindicated in:

- Known hypersensitivity to **Hyoscine Butylbromide** or any component of the formulation.
- **Narrow-angle glaucoma**.
- Prostatic hypertrophy associated with urinary retention.
- Mechanical stenosis of the gastrointestinal tract.
- Paralytic or obstructive ileus.
- Megacolon.

- Tachyarrhythmias.

It should not be used for unexplained abdominal pain without appropriate clinical evaluation.

Adverse Reactions

Common/Possible:

- Dry mouth
- Increased heart rate
- Dizziness
- Blurred vision
- Reduced sweating
- Constipation
- Difficulty in urination

Injection-related:

- Pain or irritation at the injection site
- Hypotension
- Tachycardia

Rare/Serious:

- Anaphylactic reactions
- Dyspnoea
- Severe hypotension
- Angioedema
- Urinary retention
- Acute increase in intraocular pressure in susceptible patients

Drug Interactions

The anticholinergic effects of Hyoscine Butylbromide may be increased by concomitant use with other **anticholinergic medicines**, including:

- Tricyclic antidepressants
- Tetracyclic antidepressants
- Antihistamines with anticholinergic activity
- Antipsychotic medicines with anticholinergic properties
- Other antimuscarinic agents

Concomitant use with **dopaminergic medicines** may increase gastrointestinal effects.

Other medicines that increase heart rate may potentially enhance tachycardic effects.

Patients should inform their physician about all prescription medicines, OTC medicines and herbal products being used.

Pregnancy and Lactation

Available clinical experience has not established a significant teratogenic risk; however, use during pregnancy should be based on a careful assessment of the clinical need and undertaken under medical supervision.

Hyoscine Butylbromide use during lactation should be based on medical advice because adequate data regarding its excretion into human milk are limited.

Storage

Store in a **cool, dry place**, protected from excessive heat and direct sunlight.

Keep the ampoule/vial in its original packaging until use.

Keep out of reach of children.

Do not use if the solution is discoloured, contains visible particles or if the container is damaged.

Note

For IM/slow IV use only. To be administered by or under the supervision of a qualified healthcare professional.

Butyhype Injection is intended for the short-term treatment of acute smooth-muscle spasm. Persistent or unexplained abdominal pain requires appropriate diagnostic evaluation.

Important: The final printed literature should follow the **approved product-specific Indian prescribing information**, particularly regarding exact concentration per mL, maximum daily dose, pediatric use and IV/IM administration requirements.

Manufactured in India for:

Cafoli™
LIFECARE

Cafoli Lifecare Pvt. Ltd.
(An ISO 9001: 2015 Certified Co.)
Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113
Corporate Office :
Plot no. 307, Industrial Area Phase-2,

Panchkula, Haryana -134113
TM: Trademark Applied for