Information For the Use of
Indapamide 1.5 mg Sustained Release) Tablet

Indapolite SR 1.5 Tablet

Indapolite SR 1.5 Tablet

Composition:

Each sustained-release tablet contains:

- **Indapamide** – 1.5 mg

Pharmacological Action:

Indapamide is a **thiazide-like diuretic and antihypertensive agent** that lowers blood pressure through diuretic and direct vascular effects.

- **Diuretic Action:** Reduces sodium reabsorption in the renal tubules, increasing the excretion of sodium and water.
- **Reduction in Blood Volume:** The initial diuretic effect reduces circulating fluid volume and cardiac workload.
- **Vasodilatory Action:** With continued treatment, indapamide reduces peripheral vascular resistance, contributing to sustained blood-pressure control.
- **Sustained-Release Effect:** The SR formulation provides prolonged drug release suitable for once-daily administration.

Indications:

Indapolite SR 1.5 Tablet is indicated for the management of:

- **Hypertension**
- **Essential Hypertension**
- Blood pressure control requiring once-daily diuretic therapy

---

Dosage and Administration:

- **Recommended Dose:** One tablet **once daily**, preferably in the morning, or as directed by the physician.
- **Administration:** Swallow the tablet whole with water.
- Do not crush or chew the sustained-release tablet.
- It may be taken with or without food.
- Regular monitoring of blood pressure and electrolytes may be required during treatment.

Precautions:

- Monitor **blood pressure, renal function, serum sodium, and serum potassium** periodically.
- Use cautiously in patients with **renal or hepatic impairment**.
- Indapamide may cause **hypokalemia, hyponatremia, dehydration, or hypotension**.
- Patients with diabetes should have blood glucose monitored periodically.
- Use cautiously in patients with **gout or hyperuricemia**, as indapamide may increase uric acid levels.
- Adequate fluid intake should be maintained unless fluid restriction has been prescribed.
- **Pregnancy and Lactation:** Use only when clearly indicated and under medical supervision.

Contraindications:

- Known hypersensitivity to indapamide or any component of the formulation.
- **Severe renal impairment or anuria**.
- Severe hepatic impairment associated with hepatic encephalopathy.
- Significant **hypokalemia or hyponatremia** until corrected.
- Concomitant use with medicines that may produce serious electrolyte disturbances requires careful medical assessment.

Adverse Reactions:

Common:

- Dizziness
- Headache
- Fatigue
- Increased urination
- Nausea
- Electrolyte disturbances

Less Common but Serious:

- Significant **hypokalemia or hyponatremia**
- Dehydration and hypotension
- Increased uric acid or gout
- Abnormal heart rhythm secondary to electrolyte disturbance
- Worsening renal function
- Severe hypersensitivity reactions

Patients should seek medical attention for severe weakness, confusion, fainting, persistent muscle cramps, palpitations, or marked dizziness.

Drug Interactions:

- **Lithium:** Indapamide may increase lithium concentrations and the risk of lithium toxicity.
- Other antihypertensive medicines may produce an additive blood-pressure-lowering effect.
- **NSAIDs** may reduce the antihypertensive/diuretic effect and may increase the risk of renal impairment.
- Other diuretics or medicines affecting potassium levels may increase the risk of electrolyte disturbances.
- **Digoxin:** Indapamide-induced hypokalemia may increase the risk of digoxin toxicity.
- Medicines that prolong the **QT interval** may carry increased arrhythmia risk in the presence of electrolyte abnormalities.

Storage:

- Store in a cool, dry place below **25°C**.
- Protect from moisture and direct sunlight.
- Keep the tablets in their original packaging.
- Keep out of reach of children.

Note:

This information is intended as a general prescribing guide. **Indapolite SR 1.5 Tablet should be used under the supervision of a registered medical practitioner.** The dosage and duration should be individualized according to blood pressure, renal function, electrolyte status, and clinical response.

Important: Periodic monitoring of **blood pressure, serum sodium, potassium, and renal function** is recommended during treatment, particularly in elderly patients and those receiving other diuretics or medicines affecting electrolytes.

Manufactured in India for:

Cafoli™ LIFECARE

Cafoli Lifecare Pvt. Ltd.
(An ISO 9001: 2015 Certified Co.)
Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113
Corporate Office :
Plot no. 307, Industrial Area Phase-2,

Panchkula, Haryana -134113
TM: Trademark Applied for