Information For the Use of
Isoxsuprine (10mg) Tablet

Isopiroz SR 10 Tablet

Brand Name: Isopiroz SR 10 Tablet
Dosage Form: Sustained Release Tablet
Composition:
Each sustained-release tablet contains:
• Isoxsuprine Hydrochloride IP 10 mg

Description:
Isopiroz SR 10 Tablet contains Isoxsuprine Hydrochloride, a peripheral vasodilator and beta-adrenergic agonist. It is indicated for the management of vascular insufficiency and conditions associated with impaired blood circulation. It helps in improving blood flow by relaxing vascular smooth muscles and reducing peripheral vascular resistance.

Indications:
Indicated for the treatment of:
• Peripheral vascular diseases
• Cerebral vascular insufficiency
• Raynaud's phenomenon
• Conditions associated with reduced peripheral blood flow
It helps in enhancing circulation, improving tissue perfusion, and relieving symptoms associated with vascular insufficiency.

Dosage & Administration:
Dosage should be as directed by the physician.
The tablet should be swallowed whole with water and should not be crushed or chewed.
Administration after meals may help improve gastrointestinal tolerability.

Pharmacological Action:

- **Isoxsuprine Hydrochloride directly relaxes vascular smooth muscle and stimulates beta-adrenergic receptors.**
- **It causes peripheral vasodilation, resulting in increased blood flow to affected tissues.**
- **It also reduces vascular resistance and improves microcirculation.**

These actions help in improving circulation and relieving symptoms related to vascular compromise.

Side Effects:
Common side effects include dizziness, headache, nausea, flushing, palpitations, and mild hypotension.
Rarely, allergic reactions or tachycardia may occur.

Precautions & Warnings:
• Use with caution in patients with cardiovascular disease or hypotension
• Monitor blood pressure during therapy
• Use cautiously in elderly patients

• Not recommended in patients with recent arterial bleeding
• Pregnancy and lactation use should be based on physician evaluation

---

Drug Interactions:
May interact with antihypertensive agents and other vasodilators, leading to additive hypotensive effects.
Concurrent use with sympathomimetic agents should be undertaken with caution.

Storage:
Store below 25°C in a cool, dry place.
Protect from light and moisture.
Keep out of reach of children.

Manufactured in India for:

Cafoli™
LIFECARE

Cafoli Lifecare Pvt. Ltd.
(An ISO 9001: 2015 Certified Co.)
Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113
Corporate Office :
Plot no. 307, Industrial Area Phase-2,
Panchkula, Haryana -134113
TM: Trademark Applied for