Keptoss TH 100/8 Tablet

KEPTOSS TH 100/8 TABLET

Ketoprofen (100 mg) + Thiocolchicoside (8 mg) Tablets

Composition

Each tablet contains:

- **Ketoprofen – 100 mg**
- **Thiocolchicoside – 8 mg**

Pharmacological Action

Ketoprofen is an NSAID with analgesic and anti-inflammatory properties, while thiocolchicoside is a centrally acting skeletal muscle relaxant.

It functions by:

- **Analgesic Action:** Ketoprofen reduces the production of prostaglandins involved in pain generation.
- **Anti-inflammatory Action:** Reduces inflammation, swelling and tenderness associated with musculoskeletal conditions.
- **Muscle Relaxation:** Thiocolchicoside helps reduce painful skeletal muscle spasm and excessive muscle tone.
- **Improved Mobility:** Relief of pain and muscle spasm may help improve movement and functional mobility.

Indications

Keptoss TH 100/8 Tablets may be used for the short-term management of:

- Acute painful musculoskeletal conditions associated with muscle spasm
- Acute low back pain with painful muscle spasm
- Painful spinal muscle contractures
- Musculoskeletal pain following sprains and strains
- Other acute painful conditions involving muscle spasm, as prescribed by the physician

Systemic thiocolchicoside is recommended as **short-term adjunctive treatment for painful muscle contractures associated with acute spinal conditions** in adults and adolescents aged 16 years and above.

Dosage and Administration

- **Recommended Dose:** As directed by the physician. The thiocolchicoside component should **not exceed 8 mg every 12 hours (16 mg/day)**.
- **Administration:** Swallow the tablet whole with water, preferably with or after food to help reduce gastrointestinal discomfort associated with ketoprofen.
- **Duration of Use:** Systemic oral thiocolchicoside treatment should generally be limited to **a maximum of 7 consecutive days** and should not be used for long-term treatment.
- Do not exceed the prescribed dose or duration.

Precautions

- **Pregnancy & Lactation:** Thiocolchicoside-containing systemic medicines should not be used during pregnancy or lactation. Women of childbearing potential should use effective contraception during treatment.
- **Gastrointestinal Disorders:** Use ketoprofen cautiously in patients with a history of peptic ulcer, gastritis or gastrointestinal bleeding.
- **Kidney & Liver Impairment:** Use cautiously in patients with renal or hepatic impairment; appropriate medical monitoring may be required.
- **Cardiovascular Conditions:** NSAIDs should be used cautiously in patients with hypertension, heart disease or cardiovascular risk factors.
- **Dizziness/Drowsiness:** Thiocolchicoside may cause dizziness or drowsiness in some patients. Avoid driving or operating machinery if affected.
- **Elderly Patients:** Use with appropriate medical supervision because the risk of NSAID-related gastrointestinal, renal and cardiovascular adverse effects may be increased.

Contraindications

- Known hypersensitivity to ketoprofen, thiocolchicoside, other NSAIDs or any component of the formulation.
- Active peptic ulcer or gastrointestinal bleeding.
- Previous NSAID-associated gastrointestinal bleeding or severe hypersensitivity reactions.
- Severe renal or hepatic impairment, unless specifically advised and monitored.
- **Pregnancy and lactation.**
- Women of childbearing potential who are not using adequate contraception.
- Systemic thiocolchicoside should not be used in children and adolescents below **16 years of age**.

Adverse Reactions

Common:

- Nausea
- Vomiting
- Indigestion
- Stomach discomfort or abdominal pain
- Heartburn
- Dizziness or drowsiness

- Diarrhoea

Rare but Serious:

- Gastrointestinal ulceration or bleeding
- Severe allergic reactions
- Bronchospasm, particularly in susceptible NSAID-sensitive patients
- Liver-function abnormalities
- Severe skin reactions
- Significant renal impairment

Seek immediate medical attention for gastrointestinal bleeding, severe allergic reactions, breathing difficulty or other serious symptoms.

Drug Interactions

- **Other NSAIDs:** Concurrent use with other NSAIDs may increase gastrointestinal and renal adverse effects.
- **Anticoagulants/Antiplatelet Drugs:** May increase the risk of gastrointestinal bleeding.
- **Corticosteroids:** May increase the risk of gastrointestinal ulceration and bleeding.
- **ACE inhibitors/ARBs and Diuretics:** NSAIDs may reduce their antihypertensive effect and increase renal risk in susceptible patients.
- **Alcohol:** May increase gastrointestinal irritation and other adverse effects associated with NSAIDs.
- Inform the physician about all medicines being taken before starting treatment.

Storage

- Store in a cool, dry place, preferably **below 30°C**.
- Protect from moisture and excessive heat.
- Keep the tablets in their original packaging until use.
- Keep out of reach of children.

Note

This information is intended as a **professional pharmaceutical-literature guideline** and does not replace the prescribing information approved for the specific product. **Keptoss TH 100/8 should be used under the supervision of a qualified healthcare professional.** Because systemic thiocolchicoside has specific restrictions on dose, duration, age and reproductive use, these requirements should be followed carefully.

Manufactured in India for:

Cafoli™ LIFECARE

Cafoli Lifecare Pvt. Ltd.
(An ISO 9001: 2015 Certified Co.)
Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113
Corporate Office :
Plot no. 307, Industrial Area Phase-2,
Panchkula, Haryana -134113
TM: Trademark Applied for