Information For the Use of
Ketoprofen (50mg) + Thiocolchicoside (4mg) Tablets

Keptoss TH 50/4 Tablet

KEPTOSS TH 50/4 TABLET

Thiocolchicoside (4 mg) + Ketoprofen (50 mg) Tablets

Composition

Each tablet contains:

- **Thiocolchicoside – 4 mg**
- **Ketoprofen – 50 mg**

Pharmacological Action

Ketoprofen is a non-steroidal anti-inflammatory drug (NSAID) with analgesic and anti-inflammatory activity, while thiocolchicoside is a centrally acting skeletal muscle relaxant.

It functions by:

- **Analgesic Action:** Ketoprofen reduces prostaglandin synthesis, helping relieve pain.
- **Anti-inflammatory Action:** Reduces inflammation, tenderness and swelling associated with musculoskeletal disorders.
- **Muscle Relaxation:** Thiocolchicoside helps reduce painful skeletal muscle spasm and excessive muscle tone.
- **Combined Action:** The combination provides relief from pain associated with acute musculoskeletal conditions accompanied by muscle spasm.

Indications

Keptoss TH 50/4 Tablets may be indicated for the short-term management of:

- Acute painful musculoskeletal conditions associated with muscle spasm
- Acute low back pain with painful muscle spasm
- Painful muscle contractures associated with acute spinal conditions
- Acute neck and back muscle spasm
- Painful sprains and strains associated with muscle spasm

Systemic thiocolchicoside is recommended as an adjunct for painful muscle contractures associated with acute spinal pathology in adults and adolescents **16 years and older**.

Dosage and Administration

- **Recommended Dose:** As directed by the physician. The thiocolchicoside component should not exceed **8 mg every 12 hours (16 mg/day)**.

- **Administration:** Swallow the tablet whole with water, preferably with or after food.
- **Duration of Use:** Oral systemic thiocolchicoside treatment should be limited to **a maximum of 7 consecutive days**.
- Do not exceed the prescribed dose or duration of treatment.

Precautions

- **Pregnancy & Lactation:** Thiocolchicoside should not be used during pregnancy or lactation. Women of childbearing potential should use appropriate contraception during treatment.
- **Gastrointestinal Disorders:** Use ketoprofen cautiously in patients with a history of peptic ulcer, gastritis or gastrointestinal bleeding.
- **Kidney & Liver Impairment:** Use cautiously in patients with renal or hepatic impairment under medical supervision.
- **Cardiovascular Conditions:** NSAIDs should be used cautiously in patients with hypertension, heart disease or other cardiovascular risk factors.
- **Seizure Disorders:** Thiocolchicoside should be used cautiously in patients with epilepsy or increased seizure risk.
- **Dizziness/Drowsiness:** Patients should exercise caution while driving or operating machinery if dizziness or drowsiness occurs.

Contraindications

- Known hypersensitivity to thiocolchicoside, ketoprofen, other NSAIDs or any component of the formulation.
- Active peptic ulcer or gastrointestinal bleeding.
- History of serious NSAID-related gastrointestinal reactions.
- Severe renal or hepatic impairment, where use is not considered appropriate.
- **Pregnancy and lactation.**
- Women of childbearing potential who are not using adequate contraception.
- Children and adolescents **below 16 years of age** for systemic thiocolchicoside use.

Adverse Reactions

Common:

- Nausea
- Vomiting
- Indigestion
- Abdominal discomfort
- Heartburn
- Diarrhoea
- Dizziness or drowsiness

Rare but Serious:

- Gastrointestinal ulceration or bleeding
- Severe allergic reactions

- Bronchospasm in susceptible patients
- Liver-function abnormalities
- Severe skin reactions
- Seizures or other serious neurological reactions

Seek medical attention if severe abdominal pain, gastrointestinal bleeding, breathing difficulty, severe allergic reaction or other serious symptoms occur.

Drug Interactions

- **Other NSAIDs:** May increase the risk of gastrointestinal and renal adverse effects.
- **Anticoagulants/Antiplatelet Drugs:** May increase the risk of bleeding.
- **Corticosteroids:** May increase the risk of gastrointestinal ulceration and bleeding.
- **ACE Inhibitors/ARBs and Diuretics:** NSAIDs may reduce their antihypertensive effect and increase renal risk in susceptible patients.
- **Alcohol:** May increase gastrointestinal irritation and other adverse effects associated with ketoprofen.
- Inform the physician about all medicines and supplements being taken before starting treatment.

Storage

- Store in a cool, dry place, preferably **below 30°C**.
- Protect from moisture, excessive heat and direct light.
- Keep the tablets in their original packaging until use.
- Keep out of reach of children.

Note

This information is intended as a **professional pharmaceutical-literature guideline**. **Keptoss TH 50/4 should be used under the supervision of a qualified healthcare professional.** Systemic thiocolchicoside is intended for short-term use only, with specific restrictions regarding dose, duration, age and reproductive safety.

Manufactured in India for:

Cafoli™
LIFECARE

Cafoli Lifecare Pvt. Ltd.
(An ISO 9001: 2015 Certified Co.)
Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113
Corporate Office :
Plot no. 307, Industrial Area Phase-2,
Panchkula, Haryana -134113
TM: Trademark Applied for