Information For the Use of
Latanoprost (0.005% w/v) Eye Drop

Latoshell Eye Drop

Latoshell Eye Drops

Composition:

Each mL contains:

- **Latanoprost** – 0.005% w/v

Pharmacological Action:

Latanoprost is a **prostaglandin F2α analogue** used to reduce elevated intraocular pressure (IOP).

- **Increases Aqueous Outflow:** Enhances the uveoscleral outflow of aqueous humour.
- **Reduces Intraocular Pressure:** Increased drainage of aqueous humour lowers IOP.
- **Sustained Effect:** Provides prolonged IOP reduction with once-daily administration.
- **Optic Nerve Protection:** Reduction of elevated IOP helps reduce the risk of progressive glaucomatous optic nerve damage.

Indications:

Latoshell Eye Drops is indicated for the reduction of elevated intraocular pressure in:

- **Primary Open-Angle Glaucoma**
- **Ocular Hypertension**
- Patients requiring topical IOP-lowering therapy as prescribed by an ophthalmologist

---

Dosage and Administration:

- **Recommended Dose:** Instill **1 drop into the affected eye(s) once daily**, preferably in the evening, or as directed by the ophthalmologist.
- Do not administer more than once daily, as more frequent administration may reduce the IOP-lowering effect.
- Wash hands before use.
- Tilt the head back and gently pull down the lower eyelid.
- Instill one drop without allowing the dropper tip to touch the eye or surrounding surfaces.
- Close the eye gently and apply **nasolacrimal occlusion for approximately 1–2 minutes** to reduce systemic absorption.
- If other ophthalmic medicines are prescribed, allow an appropriate interval between applications.

- Remove contact lenses before administration and wait **at least 15 minutes** before reinserting them.

Precautions:

- For **ophthalmic use only**.
- Latanoprost may gradually increase **brown pigmentation of the iris**, which can be permanent.
- It may cause increased eyelash growth and darkening of the eyelid or periocular skin.
- Use cautiously in patients with **aphakia, pseudophakia with a torn posterior lens capsule, or risk factors for macular edema**.
- Use cautiously in patients with a history of **uveitis, iritis, or herpetic keratitis**.
- Contact lenses should be removed before administration because the formulation may contain benzalkonium chloride or another preservative depending on the product.
- Avoid touching the dropper tip to the eye or other surfaces to prevent contamination.
- **Pregnancy and Lactation:** Use only when clearly indicated and under specialist supervision.

Contraindications:

- Known hypersensitivity to **latanoprost or any component of the formulation**.
- Patients with a previous serious hypersensitivity reaction to latanoprost-containing preparations.

Adverse Reactions:

Common:

- Eye redness
- Burning or stinging
- Eye irritation
- Increased eyelash growth
- Darkening of eyelid or periocular skin
- Increased iris pigmentation
- Temporary blurred vision
- Dry eye

Less Common but Serious:

- **Macular edema**
- Uveitis or iritis
- Herpetic keratitis reactivation
- Significant ocular inflammation
- Severe hypersensitivity reactions

Seek medical attention for persistent eye pain, significant visual changes, severe inflammation, or sudden deterioration of vision.

Drug Interactions:

- Concurrent use with other **prostaglandin analogues** may result in reduced IOP-lowering effectiveness or paradoxical elevation of IOP and should generally be avoided unless specifically directed.
- Other ophthalmic IOP-lowering medicines may produce an additive reduction in IOP.
- If multiple eye drops are prescribed, administer them separately with an appropriate interval.
- Inform the ophthalmologist about all prescription, OTC, and ophthalmic medicines being used.

Storage:

- **Before opening:** Store under refrigeration at **2–8°C**, protected from light, unless the approved product label specifies otherwise.
- **After opening:** Store according to the manufacturer's approved instructions and discard after the specified period.
- Do not freeze.
- Keep the bottle tightly closed.
- Keep out of reach of children.

Note:

This information is intended as a general prescribing guide. **Latoshell Eye Drops should be used under the supervision of a qualified ophthalmologist**, with periodic monitoring of intraocular pressure.

Important: Increased iris pigmentation associated with latanoprost may be **permanent**. Patients should not increase the frequency of administration beyond the prescribed once-daily dose.

Manufactured in India for:

Cafoli™ LIFECARE

Cafoli Lifecare Pvt. Ltd.
(An ISO 9001: 2015 Certified Co.)
Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113

Corporate Office :
Plot no. 307, Industrial Area Phase-2,
Panchkula, Haryana -134113
TM: Trademark Applied for