Information For the Use of
Timolol (5mg/ml) + Latanoprost (50mcg/ml) Eye Drops

Latoshell T Eye Drop

Latoshell T Eye Drops

Composition:

Each mL contains:

- **Timolol** – 5 mg
- **Latanoprost** – 50 mcg

Pharmacological Action:

Latoshell T combines a **topical beta-adrenergic blocker** with a **prostaglandin analogue** to provide complementary reduction of intraocular pressure (IOP).

- **Timolol:** Blocks beta-adrenergic receptors in the ciliary body, reducing the production of aqueous humour.
- **Latanoprost:** Increases the uveoscleral outflow of aqueous humour, resulting in reduction of IOP.
- **Combined Action:** The two complementary mechanisms provide enhanced IOP reduction compared with either agent alone in appropriately selected patients.
- **Optic Nerve Protection:** Sustained reduction of elevated IOP helps reduce the risk of progressive glaucomatous optic nerve damage.

Indications:

Latoshell T Eye Drops is indicated for the reduction of elevated intraocular pressure in:

- **Primary Open-Angle Glaucoma**
- **Ocular Hypertension**
- Patients requiring additional IOP reduction when treatment with a single agent is insufficient

---

Dosage and Administration:

- **Recommended Dose:** Instill **1 drop into the affected eye(s) once daily**, preferably at the same time each day, or as directed by the ophthalmologist.
- Do not administer more frequently than prescribed.
- Wash hands before use.
- Tilt the head back and gently pull down the lower eyelid.
- Instill one drop without allowing the dropper tip to touch the eye or surrounding surfaces.

- Close the eye gently and apply **nasolacrimal occlusion for 1–2 minutes** to reduce systemic absorption.
- If other ophthalmic medicines are being used, allow an appropriate interval between applications.
- Remove contact lenses before administration and wait **at least 15 minutes** before reinserting them.

Precautions:

- For **ophthalmic use only**.
- Timolol may be systemically absorbed and can produce **bradycardia, hypotension, bronchospasm, and cardiac conduction abnormalities**.
- Use cautiously in patients with cardiovascular disease, diabetes, thyroid disorders, or peripheral vascular disease.
- Latanoprost may cause **increased iris pigmentation**, eyelash growth, and darkening of the eyelid/periocular skin. Iris pigmentation may be permanent.
- Use cautiously in patients with a history of **uveitis, macular edema, or herpetic keratitis**.
- Latanoprost may rarely cause macular edema, particularly in patients with aphakia or pseudophakia with a damaged posterior lens capsule.
- Avoid contamination of the dropper tip.
- **Pregnancy and Lactation:** Use only when clearly indicated and under specialist supervision.

Contraindications:

- Known hypersensitivity to **timolol, latanoprost, or any component of the formulation**.
- **Bronchial asthma or a history of bronchial asthma**.
- Severe chronic obstructive pulmonary disease with significant bronchospastic disease.
- **Sinus bradycardia**.
- Second- or third-degree **atrioventricular block** without a functioning pacemaker.
- **Overt cardiac failure or cardiogenic shock**.

Adverse Reactions:

Common:

- Eye redness
- Burning or stinging
- Eye irritation
- Dry eye
- Blurred vision
- Increased eyelash growth
- Changes in iris or eyelid pigmentation

Systemic Effects:

- Bradycardia
- Hypotension
- Dizziness
- Fatigue
- Shortness of breath

Rare but Serious:

- Bronchospasm
- Cardiac conduction disturbances
- Severe bradycardia
- Macular edema
- Severe hypersensitivity reactions

Seek immediate medical attention for difficulty breathing, fainting, severe dizziness, marked reduction in heart rate, or sudden visual changes.

Drug Interactions:

- Other **topical or systemic beta-blockers** may increase beta-blocking effects.
- **Calcium-channel blockers**, cardiac glycosides, and other medicines affecting cardiac conduction may increase the risk of bradycardia or heart block.
- **Catecholamine-depleting medicines** may enhance hypotension or bradycardia.
- Other glaucoma medicines may produce additive IOP-lowering effects.
- If multiple ophthalmic preparations are prescribed, administer them separately with an appropriate interval.

Storage:

- Store according to the manufacturer's approved storage conditions, generally **2–8°C before opening** for many latanoprost-containing formulations.
- Protect from light.
- After opening, follow the **specific manufacturer's storage and discard instructions**.
- Keep the bottle tightly closed when not in use.
- Keep out of reach of children.

Note:

This information is intended as a general prescribing guide. **Latoshell T Eye Drops should be used under the supervision of a qualified ophthalmologist**, with periodic monitoring of intraocular pressure.

Important: Timolol can be systemically absorbed following ophthalmic administration. Patients with **asthma, significant COPD, bradycardia, heart block, or cardiac failure** should be carefully evaluated before treatment.

Manufactured in India for:

Cafoli Lifecare Pvt. Ltd.
(An ISO 9001: 2015 Certified Co.)
Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113
Corporate Office :
Plot no. 307, Industrial Area Phase-2,
Panchkula, Haryana -134113
TM: Trademark Applied for