Information For the Use of
Loteprednol etabonate (0.5% w/v) + Tobramycin (0.3% w/v) Eye Drop

Lotplit T Eye Drop

Lotplit T Eye Drops

Composition:

Each mL contains:

- **Loteprednol Etabonate** – 0.5% w/v
- **Tobramycin** – 0.3% w/v

Pharmacological Action:

Lotplit T combines a **topical corticosteroid** with an **aminoglycoside antibiotic** to control ocular inflammation associated with susceptible bacterial infection.

- **Anti-inflammatory Action:** Loteprednol etabonate suppresses the release and activity of inflammatory mediators, reducing ocular redness, swelling, irritation, and discomfort.
- **Antibacterial Action:** Tobramycin inhibits bacterial protein synthesis by binding to bacterial 30S ribosomal subunits, thereby inhibiting growth of susceptible bacteria.
- **Combined Action:** Provides anti-inflammatory and antibacterial activity when both are clinically indicated.

Indications:

Lotplit T Eye Drops is indicated for steroid-responsive inflammatory conditions of the eye where a **bacterial infection or risk of bacterial infection** exists, including:

- **Post-operative Ocular Inflammation**
- **Bacterial Ocular Inflammation**
- **Steroid-responsive Inflammatory Conditions associated with susceptible bacterial infection**
- Other ocular conditions as diagnosed and prescribed by an ophthalmologist

---

Dosage and Administration:

- **Recommended Dose:** Instill **1–2 drops into the affected eye(s) 4 times daily**, or as directed by the ophthalmologist.
- Shake the bottle well before use if required by the formulation.
- Wash hands before administration.
- Tilt the head back and gently pull down the lower eyelid.

- Instill the prescribed drop without touching the dropper tip to the eye, eyelids, or surrounding surfaces.
- Close the eye gently after administration.
- Apply **nasolacrimal occlusion for 1–2 minutes** when appropriate to reduce systemic absorption.
- If other ophthalmic preparations are being used, maintain an appropriate interval between applications.
- Do not use for longer than prescribed.

Precautions:

- For **ophthalmic use only**.
- Prolonged use of topical corticosteroids may cause **increased intraocular pressure, glaucoma, cataract formation, and delayed corneal healing**.
- Intraocular pressure should be monitored when treatment is prolonged or repeated.
- Corticosteroids may **mask or worsen ocular infections**, particularly fungal or viral infections.
- Use cautiously in patients with a history of **herpes simplex keratitis**, glaucoma, or corneal thinning.
- Long-term or inappropriate antibiotic use may promote **overgrowth of non-susceptible organisms**, including fungi.
- Contact lenses should generally be removed before administration and reinserted only as advised.
- Avoid touching the dropper tip to any surface to prevent contamination.
- **Pregnancy and Lactation:** Use only when clearly indicated and under medical supervision.

Contraindications:

- Known hypersensitivity to **loteprednol etabonate, tobramycin, or any component of the formulation**.
- **Acute untreated purulent ocular infections** where corticosteroid therapy may worsen the condition.
- Viral diseases of the cornea and conjunctiva, including **epithelial herpes simplex keratitis**, unless specifically directed by an ophthalmologist.
- **Fungal diseases of ocular structures**.
- Mycobacterial ocular infections.

Adverse Reactions:

Common:

- Temporary burning or stinging
- Eye irritation
- Blurred vision
- Dry eye
- Eye redness

- Discomfort following instillation

Less Common but Serious:

- Increased intraocular pressure
- Glaucoma
- Posterior subcapsular cataract with prolonged use
- Delayed corneal healing
- Secondary ocular infection
- Corneal thinning or perforation in susceptible eyes
- Severe hypersensitivity reactions

Seek prompt ophthalmic evaluation for severe eye pain, worsening redness, reduced vision, persistent discharge, or symptoms that worsen during treatment.

Drug Interactions:

- Concurrent use with other **ophthalmic corticosteroids** may increase the risk of steroid-related ocular adverse effects.
- Other topical aminoglycosides may increase the risk of local toxicity.
- Multiple ophthalmic medicines may have additive local effects; administer them separately with an appropriate interval.
- Inform the ophthalmologist about all prescription, OTC, and ophthalmic medicines being used.

Storage:

- Store in a cool, dry place below **25°C**, unless otherwise specified on the approved product label.
- Protect from direct sunlight and excessive heat.
- Keep the bottle tightly closed when not in use.
- Do not touch the dropper tip to any surface.
- Keep out of reach of children.

Note:

This information is intended as a general prescribing guide. **Lotplit T Eye Drops should be used under the supervision of a qualified ophthalmologist.** Because the product contains a corticosteroid, prolonged or unsupervised use can increase intraocular pressure and may worsen or mask certain ocular infections.

Important: If treatment continues for **10 days or longer**, intraocular pressure should generally be monitored. The combination should be used only when both **anti-inflammatory and antibacterial therapy** are clinically appropriate.

Manufactured in India for:

Cafoli Lifecare Pvt. Ltd.
(An ISO 9001: 2015 Certified Co.)
Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113
Corporate Office :
Plot no. 307, Industrial Area Phase-2,
Panchkula, Haryana -134113
TM: Trademark Applied for