Information For the Use of
Mebeverine (200mg) SR Capsules

Mebihusk SR 200 Capsule

Mebihusk SR 200 Capsule

Composition:

Each sustained-release capsule contains:

- **Mebeverine** – 200 mg

Pharmacological Action:

Mebeverine is a **direct-acting antispasmodic** that acts primarily on the smooth muscles of the gastrointestinal tract.

- **Antispasmodic Action:** Relaxes intestinal smooth muscle and reduces painful gastrointestinal spasms.
- **Smooth Muscle Regulation:** Helps relieve abnormal intestinal contractions without significantly affecting normal intestinal motility.
- **Visceral Spasm Relief:** Reduces abdominal cramping and discomfort associated with intestinal spasm.
- **Local Gastrointestinal Action:** Primarily exerts its therapeutic effect within the gastrointestinal tract.

Indications:

Mebihusk SR 200 Capsule is indicated for the symptomatic management of:

- **Irritable Bowel Syndrome (IBS)**
- **Abdominal Cramps and Spasms**
- **Functional Bowel Disorders**
- **Intestinal Spasm**
- **Abdominal Pain and Discomfort associated with gastrointestinal spasm**

Dosage and Administration:

- **Recommended Dose:** One capsule **twice daily** or as directed by the physician.
- **Administration:** Take approximately **20 minutes before meals** with sufficient water.
- Swallow the sustained-release capsule whole.
- **Do not crush or chew** the capsule contents unless specifically permitted by the approved formulation.

- Treatment duration should be determined according to the patient's symptoms and clinical response.

Precautions:

- Use cautiously in patients with severe or unexplained abdominal pain.
- Persistent abdominal pain accompanied by **fever, vomiting, blood in stools, marked abdominal tenderness, or unexplained weight loss** requires medical evaluation.
- Mebeverine is intended for symptomatic relief and does not treat underlying structural gastrointestinal disease.
- Patients with severe hepatic or renal impairment should consult a physician before use.
- **Pregnancy and Lactation:** Use only when clearly indicated and under medical supervision.
- If symptoms persist or worsen despite treatment, medical evaluation is recommended.

Contraindications:

- Known hypersensitivity to **mebeverine** or any component of the formulation.
- Use in children should follow the age-specific recommendations of the approved product information.
- Patients with suspected intestinal obstruction should undergo appropriate medical evaluation rather than relying on antispasmodic treatment.

Adverse Reactions:

Common:

- Nausea
- Indigestion
- Headache
- Mild dizziness

Rare but Serious:

- Hypersensitivity reactions
- Skin rash
- Urticaria
- Angioedema
- Severe allergic reactions, including anaphylaxis

Patients should discontinue treatment and seek medical attention if facial swelling, breathing difficulty, or other severe allergic symptoms occur.

Drug Interactions:

- Mebeverine has **no major clinically significant drug interactions** commonly reported.
- Inform the physician about all prescription medicines, OTC medicines, and herbal supplements being used.
- Avoid combining multiple antispasmodic medicines without medical advice.

Storage:

- Store in a cool, dry place below **25°C**.
- Protect from moisture and direct sunlight.
- Keep the capsule in its original packaging.
- Keep out of reach of children.

Note:

This information is intended as a general prescribing guide. Mebihusk SR 200 Capsule should be used according to the directions of a registered medical practitioner. The sustained-release formulation should be swallowed whole and not crushed or chewed.

Persistent, severe, or unexplained abdominal symptoms should be medically evaluated rather than managed solely with an antispasmodic.

Manufactured in India for:

Cafoli™ LIFECARE

Cafoli Lifecare Pvt. Ltd.
(An ISO 9001: 2015 Certified Co.)
Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113
Corporate Office :
Plot no. 307, Industrial Area Phase-2,
Panchkula, Haryana -134113
TM: Trademark Applied for