# Melaturn 3 MD Tablet

## MELATURN 3 MD

### Composition

Each Mouth Dissolving Tablet contains:

**Melatonin 3 mg**
Excipients … q.s.

### Pharmacological Action

**Melatonin** is a naturally occurring hormone primarily secreted by the pineal gland. It plays an important role in regulating the body's **sleep-wake cycle and circadian rhythm**.

Exogenous melatonin acts on melatonin receptors involved in synchronization of the biological clock. Administration at the appropriate time can help regulate disturbed sleep timing and facilitate normalisation of the sleep-wake rhythm.

### Indications

Melaturn 3 MD may be indicated for the symptomatic management of:

- Disturbed sleep-wake rhythm
- Circadian rhythm disturbances
- Short-term sleep disturbance associated with jet lag
- Difficulty initiating sleep associated with altered sleep timing

For the exact marketed indication, the approved product-specific prescribing information should be followed. A 3 mg melatonin formulation is established internationally for short-term treatment of jet lag in adults.

### Dosage and Administration

**Adults:**
The usual dose of immediate-release melatonin is **3 mg once daily**, taken at the appropriate bedtime according to the indication. For jet lag, the dose is taken at the usual bedtime at the destination, generally for a short duration.

**Administration:**
As this is a **Mouth Dissolving Tablet (MD)**, place the tablet in the mouth and allow it to disintegrate before swallowing, according to the product's approved administration instructions.

Use the **lowest effective dose for the shortest duration** appropriate to the indication.

## Precautions

- May cause **drowsiness or reduced alertness**; patients should avoid driving or operating machinery if affected.
- Use cautiously in patients with a history of **seizures/epilepsy**, as melatonin may increase seizure frequency in susceptible individuals.
- Use with caution in patients with **autoimmune disorders**.
- Caution is advised in patients with significant **hepatic or renal impairment**.
- Patients with diabetes or impaired glucose tolerance should use melatonin under medical supervision.
- Alcohol should preferably be avoided because it may impair sleep and increase drowsiness.
- The timing of administration is important because inappropriate timing may reduce the desired circadian effect.

## Contraindications

Melaturn 3 MD is contraindicated in patients with:

- Hypersensitivity to melatonin or any component of the formulation.
- Any other contraindication specified in the approved product-specific prescribing information.

## Adverse Reactions

**Common/possible:**

- Headache
- Drowsiness
- Dizziness
- Nausea
- Fatigue or lethargy

**Uncommon/rare:**

- Irritability or nervousness
- Abnormal dreams or nightmares
- Anxiety or restlessness
- Dry mouth
- Abdominal discomfort
- Changes in blood pressure
- Hypersensitivity reactions

Drowsiness and headache are among the commonly reported adverse effects with melatonin formulations.

## Drug Interactions

Melatonin may interact with:

- **Fluvoxamine**, which can substantially increase melatonin exposure.
- Other medicines producing **sedation/CNS depression**, potentially increasing drowsiness.
- **Anticoagulants/antiplatelet medicines**, where increased bleeding-related effects may require monitoring.
- Medicines affecting hepatic metabolism of melatonin.
- Alcohol, which may increase impairment and adversely affect sleep.

Patients should inform their healthcare professional about all prescription, OTC and herbal medicines being used.

## Pregnancy and Lactation

Melatonin is **not recommended during pregnancy or breastfeeding** unless specifically advised by a qualified healthcare professional, because adequate safety information is not established.

## Storage

Store in a **cool, dry place**, protected from light and moisture.

Keep the tablets in their original packaging and **out of reach of children**.

## Note

**For use under medical supervision.**

Melatonin should be taken at the appropriate time according to the intended indication. Patients experiencing persistent or recurrent sleep problems should undergo appropriate medical evaluation rather than continuing self-treatment indefinitely.

Manufactured in India for:

Cafoli™
LIFECARE

**Cafoli Lifecare Pvt. Ltd.**
**(An ISO 9001: 2015 Certified Co.)**
Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113
**Corporate Office :**
Plot no. 307, Industrial Area Phase-2,
Panchkula, Haryana -134113

TM: Trademark Applied for