Melaturn 5 MD Tablet

MELATURN 5 MD TABLETS

Composition

Each Mouth Dissolving Tablet contains:

Melatonin 5 mg
Excipients … q.s.

Pharmacological Action

Melatonin is a naturally occurring hormone produced primarily by the pineal gland. It plays an important role in regulating the body's **sleep-wake cycle and circadian rhythm**.

Exogenous melatonin acts on melatonin receptors involved in synchronisation of the biological clock. When administered at an appropriate time, it may facilitate sleep initiation and help regulate disturbed sleep-wake patterns.

Indications

Melaturn 5 MD Tablets may be used for the management of:

- Sleep-onset difficulties associated with disturbed sleep patterns
- Circadian rhythm disturbances
- Jet-lag-associated sleep disturbance
- Altered sleep-wake cycle

The exact indication should correspond to the approved product-specific prescribing information.

Dosage and Administration

Adults:
The dose should be individualized according to the indication and response. **Melaturn 5 MD should be taken only in the dose recommended by the physician.**

For certain approved melatonin products, **5 mg once daily** is used for short-term jet-lag treatment when a 3 mg dose is inadequate; the exact dose and duration depend on the approved indication.

Administration:
Place the mouth-dissolving tablet on the tongue and allow it to disintegrate before swallowing, according to the product's approved administration instructions.

The timing of administration is important because inappropriate timing may reduce the desired circadian effect.

Precautions

- May cause **drowsiness, dizziness or reduced alertness**.
- Avoid driving or operating machinery if affected by drowsiness.
- Avoid alcohol because it may impair sleep and reduce the beneficial effects of melatonin.
- Use cautiously in patients with **epilepsy or a history of seizures**.
- Use cautiously in patients with autoimmune disorders.
- Patients with hepatic impairment or severe renal impairment should use melatonin only under medical supervision.
- In patients with diabetes or impaired glucose tolerance, timing around meals should be considered because melatonin may affect glucose control.
- Do not combine with other sleep-inducing or sedative medicines without medical advice.

Contraindications

Melaturn 5 MD Tablets are contraindicated in patients with:

- Hypersensitivity to melatonin or any component of the formulation.
- Any additional contraindication specified in the approved product-specific prescribing information.

Adverse Reactions

Common/possible:

- Drowsiness
- Headache
- Dizziness
- Fatigue
- Nausea
- Lethargy

Other possible reactions:

- Abnormal dreams or nightmares
- Irritability
- Restlessness
- Anxiety
- Dry mouth
- Gastrointestinal discomfort
- Changes in blood pressure

Patients experiencing persistent or troublesome adverse effects should consult their physician.

Drug Interactions

Melatonin may interact with:

- **Fluvoxamine** and other CYP1A2 inhibitors, which may increase melatonin concentrations.
- CYP1A2 inducers such as **carbamazepine and rifampicin**, which may reduce melatonin exposure.
- Sedatives, hypnotics and other **CNS depressants**, which may increase drowsiness.
- **Anticoagulants**, including warfarin and some direct-acting oral anticoagulants, where increased anticoagulant effects may require clinical monitoring.
- Certain antihypertensive medicines, including nifedipine.
- Alcohol, which may impair sleep and increase impairment.

Patients should inform their physician about all prescription, OTC and herbal medicines being taken.

Pregnancy and Lactation

Melatonin should generally **not be used during pregnancy or breastfeeding unless specifically recommended by a healthcare professional**, as adequate safety information is limited.

Storage

Store in a **cool, dry place**, protected from light and moisture.

Keep the tablets in their original packaging and **out of reach of children**.

Note

For use under medical supervision.

Do not exceed the recommended dose. Avoid combining with other melatonin-containing or sedative preparations unless advised by a healthcare professional. Persistent or recurrent sleep problems require appropriate medical evaluation.

Manufactured in India for:

Cafoli™
LIFECARE

Cafoli Lifecare Pvt. Ltd.
(An ISO 9001: 2015 Certified Co.)
Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113

Corporate Office :
Plot no. 307, Industrial Area Phase-2,
Panchkula, Haryana -134113
TM: Trademark Applied for