Information For the Use of
Amlodipine (5mg) + Metoprolol Succinate (50mg) Tablet

Amdimet 50 Tablet

Amdimet 50 Tablets

Composition:

Each tablet contains:

- **Metoprolol Succinate IP** – 50 mg
- **Amlodipine IP** – 5 mg

Pharmacological Action:

Amdimet 50 Tablets combines a cardioselective beta-blocker with a long-acting calcium channel blocker to provide complementary cardiovascular and antihypertensive effects.

- **Beta-1 Blockade:** Metoprolol selectively blocks β1-adrenergic receptors, reducing heart rate, myocardial contractility, and cardiac workload.
- **Calcium Channel Blockade:** Amlodipine inhibits calcium influx into vascular smooth muscle, producing prolonged arterial vasodilation and reducing peripheral vascular resistance.
- **Antihypertensive Action:** The combination lowers blood pressure through reduction of cardiac output and peripheral vascular resistance.
- **Antianginal Action:** Reduction in myocardial oxygen demand together with coronary and peripheral vasodilation may help relieve anginal symptoms in appropriate patients.

Indications:

Amdimet 50 Tablets is indicated for the management of:

- **Hypertension**
- **Hypertension inadequately controlled with monotherapy**
- **Stable Angina Pectoris**
- **Patients requiring combination cardiovascular therapy**, as determined by the physician

---

Dosage and Administration:

- **Recommended Dose:** One tablet once daily or as directed by the physician.
- **Administration:** Swallow the tablet whole with water. It may be taken with or immediately after food.
- Take the tablet at approximately the same time each day.
- If the formulation is modified-release, do not crush or chew the tablet.

- Do not discontinue metoprolol abruptly without medical advice.

Precautions:

- Monitor **blood pressure and heart rate** regularly.
- Use cautiously in patients with **bradycardia, heart block, peripheral vascular disease, or cardiac dysfunction**.
- Metoprolol may mask some symptoms of **hypoglycemia** in diabetic patients.
- Use cautiously in patients with **asthma or bronchospastic disease**.
- Amlodipine may cause **peripheral edema**; persistent or severe swelling should be reported.
- Careful dose selection may be required in patients with hepatic impairment.
- **Pregnancy and Lactation:** Use only when clearly indicated and under medical supervision.

Contraindications:

- Known hypersensitivity to metoprolol, amlodipine, or any component of the formulation.
- Severe bradycardia.
- Second- or third-degree AV block without a functioning pacemaker.
- Sick sinus syndrome without a functioning pacemaker.
- Cardiogenic shock.
- Decompensated heart failure.
- Severe hypotension.

Adverse Reactions:

Common:

- Dizziness
- Fatigue
- Headache
- Bradycardia
- Hypotension
- Ankle or peripheral edema
- Flushing
- Nausea

Less Common but Serious:

- Significant bradycardia
- Severe hypotension
- Worsening heart failure
- Bronchospasm

- Conduction abnormalities
- Severe hypersensitivity reactions

Drug Interactions:

- Other **beta-blockers** or heart-rate-lowering medicines may increase the risk of bradycardia.
- **Verapamil and diltiazem** may further slow cardiac conduction and heart rate.
- Other antihypertensive medicines may produce additive blood-pressure-lowering effects.
- Certain CYP2D6 inhibitors may increase metoprolol exposure.
- CYP3A4 inhibitors may increase amlodipine exposure.
- Antidiabetic medicines may require monitoring because metoprolol can mask some symptoms of hypoglycemia.

Storage:

- Store in a cool, dry place below **25°C**.
- Protect from moisture and direct sunlight.
- Keep out of reach of children.

Note:

This information is intended as a general prescribing guide. Amdimet 50 Tablets should be used under the supervision of a registered medical practitioner. The dosage and duration of treatment should be individualized according to the patient's blood pressure, heart rate, cardiovascular condition, and clinical response.

Manufactured in India for:

Cafoli™ LIFECARE

Cafoli Lifecare Pvt. Ltd.
(An ISO 9001: 2015 Certified Co.)
Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113
Corporate Office :
Plot no. 307, Industrial Area Phase-2,
Panchkula, Haryana -134113
TM: Trademark Applied for